

BÖLÜM

8

# KARACİĞER KİTLELERİ

*Ayşe Gül ALIMLI<sup>1</sup>*

**Vaka 1:** Hepatoblastom

**Vaka 2:** Fibrolamellar Hepatosellüler Karsinom

**Vaka 3:** Metastatik İnfiltratif Nöroblastom

**Vaka 4:** Metastatik Lenfoma

**Vaka 5:** Hepatik Hemanjiom

**Vaka 6:** Hepatik Adenomatozis

**Vaka 7:** Fokal Nodüler Hiperplazi (FNH)

**Vaka 8:** Karaciğer Kist Hidatik

---

<sup>1</sup> Pediatrik Radyoloji Uzmanı, Ankara Şehir Hastanesi Çocuk Hastanesi Çocuk Radyolojisi, drsahin80@hotmail.com

## Tuzaklar

Tip 5 KH solid kitle ve tip 1 KH basit kist olarak değerlendirilmemelidir.

## Tedavi ve yaklaşım

Albendazolla tek başına medikal tedavi seçeneği olabileceği gibi, albendazolla birlikte perkütan girişimsel tedavi veya cerrahi tedavi de kullanılabilir (22).

## KAYNAKÇA

- Shelmerdine SC, Roebuck DJ, Towbin AJ, et al. MRI of pediatric liver tumours: How we review and report. *Cancer Imaging*. 2016;16(1):21. doi: 10.1186/s40644-016-0083-3.
- Jha P, Chawla SC, Tavri S, et al. Pediatric liver tumors--a pictorial review. *Eur Radiol*. 2009;19(1):209-219. doi: 10.1007/s00330-008-1106-7.
- Chavhan GB, Siddiqui I, Ingley KM, et al. Rare malignant liver tumors in children. *Pediatr Radiol*. 2019;49(11):1404-1421. doi: 10.1007/s00247-019-04402-8.
- Baheti AD, Luana SA, Li N, et al. Contrast-enhanced CT features of hepatoblastoma: Can we predict histopathology? *Clin Imaging* 2017;44:33-37
- Chung EM, Lattin GE, Cube R, et al. From the archives of the AFIP: Pediatric liver masses: radiologic-pathologic correlation. Part 2. Malignant tumors. *Radiographics*. 2011;31:483-507
- Towbin AJ, Meyers RL, Woodley H, et al. 2017 PRETEXT: radiologic staging system for primary hepatic malignancies of childhood revised for the Pediatric Hepatic International Tumour Trial (PHITT). *Pediatr Radiol*. 2018;48(4):536-554. doi: 10.1007/s00247-018-4078-z.
- Fernandez-Pineda I, Cabello-Laureano R. Differential diagnosis and management of liver tumors in infants. *World J Hepatol*. 2014 27;6(7):486-495. doi: 10.4254/wjh.v6.i7.486.
- Kelly D, Sharif K, Brown RM, et al. Hepatocellular carcinoma in children. *Clin Liver Dis*. 2015;19(2):433-447. doi: 10.1016/j.cld.2015.01.010.
- Chaturvedi A, Katzman PJ, Franco A. Neonatal neuroblastoma 4s with diffuse liver metastases (Pepper syndrome) without an adrenal/extraadrenal primary identified on imaging. *J Radiol Case Rep*. 2018;31:12(3):18-27. doi:10.3941/jrcr.v12i3.3300.
- Hegde SV, Dillman JR, Lopez MJ, et al. Imaging of multifocal liver lesions in children and adolescents. *Cancer Imaging*. 2013;12:516-529. doi: 10.1102/1470-7330.2012.0045.
- Karacaaltincaba M, Haliloglu M, Akpınar E, Akata D, Özmen M, Ariyurek M, Akhan O. Multidetector CT and MRI findings in periportal space pathologies. *Eur J Radiol*. 2007;61(1):3-10.
- Ernst L, Grabhorn E, Brinkert F, et al. Infantile Hepatic Hemangioma: Avoiding Unnecessary Invasive Procedures. *Pediatr Gastroenterol Hepatol Nutr*. 2020;23(1):72-78. doi: 10.5223/pghn.2020.23.1.72.
- Regier TS, Ramji FG. Pediatric hepatic hemangioma. *Radiographics*. 2004;24(6):1719-1724.
- Chiorean L, Cui XW, Tannapfel A, et al. Benign liver tumors in pediatric patients Review with emphasis on imaging features. *World J Gastroenterol*. 2015;28(21):8541-8561. doi: 10.3748/wjg.v21.i28.8541.
- Radhi JM, Loewy J. Hepatocellular adenomatosis associated with hereditary haemochromatosis. *Postgrad Med J*. 2000;76(892):100-102.
- Cannon RO, Dusheiko GM, Long JA, et al. Hepatocellular adenoma in a young woman with beta-thalassemia and secondary iron overload. *Gastroenterology*. 1981;81(2):352-355.
- Shuangshoti S, Thaicharoen A. Hepatocellular adenoma in a beta-thalassemic woman having secondary iron overload. *J Med Assoc Thai*. 1994;77(2):108-112.
- Patacsil SJ, Noor M, Leyva A. A Review of Benign Hepatic Tumors and Their Imaging Characteristics. *Cureus*. 2020;29:12(1):6813. doi: 10.7759/cureus.6813.
- Nie P, Yang G, Guo J, et al. A CT-based radiomics nomogram for differentiation of focal nodular hyperplasia from hepatocellular carcinoma in the non-cirrhotic liver. *Cancer Imaging*. 2020;24:20(1):20. doi: 10.1186/s40644-020-00297-z.
- Zarfati A, Chambers G, Pio L, et al. Management of focal nodular hyperplasia of the liver: Experience of 50 pediatric patients in a tertiary center. *J Pediatr Surg*. 2020 Jan 29. pii: S0022-3468(20)30045-2. doi:10.1016/j.jpedsurg.2020.01.009.
- Dietrich CF, Douira-Khomsy W, Gharbi H, et al. Cystic and alveolar echinococcosis of the hepatobiliary tract - the role of new imaging techniques for improved diagnosis. *Med Ultrason*. 2020;1;22(1):75-84. doi: 10.11152/mu-2421.
- Eyüboğlu TŞ, Gürsoy TR, Aslan AT, et al. Ten-year follow-up of children with hydatid cysts. *Turk Pediatri Ars*. 2019;25;54(3):173-178. doi: 10.14744/TurkPediatriArs.2019.24119. eCollection 2019.
- Turgut AT, Akhan O, Bhatt S, et al. Sonographic spectrum of hydatid disease. *Ultrasound Q*. 2008;24(1):17-29. doi: 10.1097/RUQ.0b013e318168f0d1.
- Gharbi HA, Hassine W, Brauner MW, et al. Ultrasound examination of the hydatid liver. *Radiology*. 1981;139:459-463.
- WHO Informal Working Group. International classification of ultrasound images in cystic echinococcosis for application in clinical and field epidemiological settings. *Acta Trop*. 2003;85:253-261.