## CHAPTER 20

## RAPID HOSPITAL DISCHARGE IN VASCULAR SURGERY

Tolga TOTOZ1

## INTRODUCTION

There have been many advances in medicine in the last century, especially in the surgical branches, with technological developments. In parallel with technological and economic developments, the proportion of elderly people in society has also increased. This situation has led to encountering a patient group whose comorbidity has increased even more in medicine. As a result, prolonged hospitalization and more advanced technological devices have also increased health expenditures. In order to reduce health costs, health protocols have been implemented to ensure rapid discharge of patients since the second half of the 1990's (1). The primary purpose of these protocols is to increase patient safety and comfort and reduce the cost by shortening the discharge process (1). The answer to the questions "How can we ensure that surgical patients are discharged as soon as possible?" or "What are the conditions that will quickly ensure well-being?" can be provided by applying these protocols. Later, it was thought that systems that could benefit from a multidisciplinary approach in terms of resources and costs in different surgical applications should be revealed, and Enhanced Recovery After Surgery (ERAS) protocols were developed for this purpose. A consensus protocol for patients undergoing colon surgery

 $<sup>^1\,</sup>$  MD. Department of Anesthesiology and Reanimation Nişantaşı University Vocational High School, tolgatotoz@hotmail.com.

## REFERENCES

- 1. Engelman RM, Rousou JA, Flack JE, Deaton DW, Humphrey CB, Ellison LH, et al. Fast-track recovery of the coronary bypass patient. Ann Thorac Surg. 1994; 58(6): 1742-6.
- 2. Fearon KCH, Ljungqvist O, Von Meyenfeldt M, Revhaug A, Dejong CHC, Lassen K, et al. Enhanced recovery after surgery: a consensus review of clinical care for patients undergoing colonic resection. Clin Nutr. 2005 Jun; 24(3): 466-77.
- 3. Ljungqvist O, Scott M, Fearon KC. Enhanced recovery after surgery: A Review. JAMA Surg. 2017; 152(3): 292-8.
- 4. ERAS Türkiye Derneği. "ERAS protokollerinin temel ögeleri". Erişim: 22.02.2022. https://eras.org.tr.
- 5. Stammers AN, Kehler DS, Afilalo J, Avery LJ, Bagshaw SM, Grocott HP, et al. Protocol for the PREHAB study- Pre-operative Rehabilitation for Reduction of Hospitalization After Coronary Bypass and Valvular Surgery: a randomised controlled trial. BMJ Open. 2015; 5(3): e007250.
- 6. Waite I, Deshpande R, Baghai M, Massey T, Wendler O, Greenwood S. Homebased preoperative rehabilitation (prehab) to improve physical function and reduce hospital length of stay for frail patients undergoing coronary artery bypass graft and valve surgery. J Cardiothorac Surg. 2017; 12(1): 91.
- 7. Valkenet K, Trappenburg JCA, Hulzebos EH, Meeteren NLU, Backx FJG. Effects of a pre-operative home-based inspiratory muscle training programme on perceived health-related quality of life in patients undergoing coronary artery bypass graft surgery. Physiotherapy. 2017 Sep; 103(3): 276-82.
- 8. Laohachai K, Winlaw D, Selvadurai H, Gnanappa GK, d'Udekem Y, Celermajer Det, et al. Inspiratory muscle training is associated with improved inspiratory muscle strength, resting cardiac output, and the ventilatory efficiency of excercise in patients with a fontan circulation. J Am Heart Assoc. 2017; 21: 6(8): e 005750.
- 9. Planas A, Clara A, Marrugat J, Pou JM, Gasol A, de Moner A, et al. Age at onset of smoking is an independent riskfactor in peripheral artery disease development. J Vasc Surg. 2002; 35: 506-9.
- 10. Ogawa M, Izawa KP, Satomi-Kobayashi S, Kitamura A, Ono R, Sakai Y, et al. Poor preoperative nutritional status is an important predictor of the retardation of rehabilitation after cardiac surgery in elderly cardiac patients. Aging Clin Exp Res. 2017; 29(2): 283-90.
- 11. Jie B, Jiang ZM, Nolan MT, Zhu SN, Yu K, Kondrup J. Impact of preoperative nutritional support on clinical outcome in abdominal surgical patients at nutritional risk. Nutrition. 2012;28(10):1022-1027.
- 12. Yu PJ, Cassiere HA, Dellis SL, Manetta F, Kohn N, Hartman AR. Impact of preoperative prealbumin on outcomes after cardiac surgery. JPEN J Parenter Enteral Nutr. 2015; 39(7): 870-4.

- 13. Ljungqvist O. Modulating postoperative insulin resistance by preoperative carbohydrate loading. Best Pract Res Clin Anaesthesiol. 2009; 23(4): 401-9.
- 14. Wong J, Zoungas S, Wright C, Teede H. Evidence-based guidelines for perioperative management of diabetes in cardiac and vascular surgery. World J Surg. 2010; 34(3): 500-3.
- 15. Bode LG, Kluytmans JA, Wertheim HF, Bogaers D, Vandenbroucke-Grauls CM, Roosendaal R, et al. Preventing surgical-site infections in nasal carriers of Staphylococcus aureus. N Engl J Med. 2010; 362(1): 9-17.
- 16. Edwards FH, Engelman RM, Houck P, Shahian DM, Bridges CR; Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: antibiotic prophylaxis in cardiac surgery, part I: duration. Ann Thorac Surg. 2006; 81(1): 397-404.
- 17. Engelman R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, et al; Workforce on Evidence-Based Medicine, Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: antibiotic prophylaxis in cardiac surgery, part II: antibiotic choice. Ann Thorac Surg. 2007; 83 (4): 1569-76.
- 18. Lavallée JF, Gray TA, Dumville J, Russell W, Cullum N. The effects of care bundles on patient outcomes: a systematic review and meta-analysis. Implement Sci. 2017; 12(1): 142.
- 19. Mutters NT, De Angelis G, Restuccia G, Di Muzio F, Schouten J, Hulscher M, et al. Use of evidence-based recommendations in an antibiotic care bundle for the intensive care unit. Int J Antimicrob Agents. 2018; 51(1): 65-70.
- 20. Ozyaprak B, Kahraman N, Erkan G, Eroglu A. Experience of Peripheral Arterial Disease Surgery with Tumescent Local Anesthesia. Int J AnesthRes. 2019; 7(3): 572-6
- 21. Sayan HE. Varis cerrahisinde anestezi. Kahraman N, editör. Venöz ve Lenfatik Sistem Hastalıkları.1. Baskı. Ankara: Akademisyen Kitabevi; 2020. p.233-46.
- 22. Osawa EA, Rhodes A, Landoni G, Galas FR, Fukushima JT, Park CH, et al. Effect of perioperative goal-directed hemodynamic resuscitation therapy on outcomes following cardiac surgery: a randomized clinical trial and systematic review. Crit Care Med. 2016; 44(4):724-33.
- 23. Sun Y, Chai F, Pan C, Romeiser JL, Gan TJ. Effect of perioperative goal-directed hemodynamic therapy on postoperative recovery following major abdominal surgery a systematic review and meta-analysis of randomized controlled trials. Crit Care. 2017; 21(1): 141.
- 24. Frank MS, Tran MK, Fleisher Lee A, Elrahmany KH. Clinical importance of body temparature in the surgecal patient. Journal of Thermal Biology. 2000; 25: 151-5.
- 25. Türk Anesteziyoloji Derneği (TARD) Anestezi Kılavuzları. İstemeyen perioperatif hipoterminin önlenmesi rehberi. TARD/K2013/1. Erişim: 04 Şubat 2021. https://www.tard.org.tr.
- 26. Wick EC, Grant MC, Wu CL. Postoperative multimodal analgesia pain management with nonopioid analgesics and techniques: a review. JAMA Surg. 2017; 152(7): 691-7.

- 27. Apfel CC, Turan A, Souza K, Pergolizzi J, Hornuss C. Intravenous acetaminophen reduces postoperative nausea and vomiting: a systematic review and meta-analysis. Pain. 2013; 154(5): 677-89.
- 28. Çelebi N, Soylu E, Yıldırım MD, Durmuş T, Varol H, Alkan A. Ortognatik cerrahi hastalarında postoperatif kusma ve bulantının önlenmesi. Atatürk Üniv Diş Hek Fak. Derg. 2013; 21(3): 362-5.
- 29. Menda F, Köner O, Sayın M, Ergenoğlu M, Küçükaksu S, Aykaç B. Effects of singledose gabapentin on postoperative pain and morphine consumption after cardiac surgery.J Cardiothorac Vasc Anesth. 2010; 24(5): 808-13.
- 30. Borde DP, Futane SS, Asegaonkar B, Apsingekar P, Khade S, Khodve B, et al. Effect of perioperative pregabalin on postoperative quality of recovery in patients undergoing off-pump coronary artery bypass grafting (OPCABG): a prospective, randomized, double-blind trial. J Cardiothorac Vasc Anesth. 2017; 31(4): 1241-5.
- 31. Özyaprak B, Kahraman N, Anarat K, Gamlı M, Erkan G, Eroğlu A. Pain Management in Patients with Peripheral Arterial Disease CBU-SBED. 2020; 7(1):12 -7.