

# Ev Hemodiyalizi Komplikasyonları

# 8

Tushar Chopra, Lakshmi Kannan ve Emaad M. Abdel-Rahman

Çeviri:

Prof Dr. Ülver Derici, Uzm. Dr. Zeynep Ural

## ANA BAŞLIKLAR

### GENEL DEĞERLENDİRİMLER

### TEKNİK KOMPLİKASYONLAR

Kanama

Hava Embolisi

Su Sistemi Problemleri

### MEDİKAL KOMPLİKASYONLAR

Hipotansiyon

Giriş Yolu

Giriş Yolu İlişkili Olaylar

Giriş Yolu İlişkili Enfeksiyonlar

Rezidüel Renal Fonksiyonun Kaybı

Elektrolit Anormallilikleri

Beslenme

Uyku

### PSİKOSOSYAL KOMPLİKASYONLAR

Uyumsuzluk

Tedavi Uyumsuzluğu

Bakıcıının Tükenmişliği/Bırakması

İstihdam Sorunları

Amerika Birleşik Devletleri’nde Son Dönem Böbrek Hastalığı (SDBH) ortalaması prevalansı milyonda 2160,7'dir. Bu hastaların sadece %2'si ev hemodiyalizi (EHD) yapmaktadır.<sup>1</sup> EHD hastaları son yıllarda giderek artmakta ve

işverenleriyle iletişim kurmalı ve gerçekçi bekentilere sahip olmalıdır. Diyaliz ünitesinde çalışan sosyal hizmet uzmanları, hastaların diyaliz tedavileri ve işlerini dengede tutma yollarını bulmalarına yardımcı olabilir. Gece EHD'ye geçmek bazı hastalar için bir çözüm olabilir.

## KAYNAKLAR

1. Saran R, R Bruce, Abbott KC, et al. US Renal Data System 2018 Annual Data Report: epidemiology of kidney disease in the United States. *Am J Kidney Dis.* 2019;73(3 suppl 1):A7-A8.
2. Cullerton BF, Walsh M, Klarenbach SW, et al. Effect of frequent nocturnal hemodialysis vs conventional hemodialysis on left ventricular mass and quality of life: a randomized controlled trial. *JAMA.* 2007;298(11):1291-1299.
3. Hall YN, Larive B, Painter P, et al. Effects of six versus three times per week hemodialysis on physical performance, health, and functioning: Frequent Hemodialysis Network (FHN) randomized trials. *Clin J Am Soc Nephrol.* 2012;7(5):782-794.
4. Hladunewich MA, Hou S, Odutayo A, et al. Intensive hemodialysis associates with improved pregnancy outcomes: a Canadian and United States cohort comparison. *J Am Soc Nephrol.* 2014;25(5):1103-1109.
5. Lockridge RS, Kjellstrand CM. Nightly home hemodialysis: outcome and factors associated with survival. *Hemodial Int.* 2011;15(2):211-218.
6. Marshall MR, van der Schriek N, Lilley D, et al. Independent community house hemodialysis as a novel dialysis setting: an observational cohort study. *Am J Kidney Dis.* 2013;61(4):598-607.
7. Weinhandl ED, Liu J, Gilbertson DT, Arneson TJ, Collins AJ. Survival in daily home hemodialysis and matched thrice-weekly in-center hemodialysis patients. *J Am Soc Nephrol.* 2012;23(5):895-904.
8. Tennankore KK, d'Gama C, Faraturo R, Fung S, Wong E, Chan CT. Adverse technical events in home hemodialysis. *Am J Kidney Dis.* 2015;65(1):116-121.
9. Wong B, Zimmerman D, Reintjes F, et al. Procedure-related serious adverse events among home hemodialysis patients: a quality assurance perspective. *Am J Kidney Dis.* 2014;63(2):251-258.
10. Kennedy C, McGrath-Chong M, Arustei D, et al. A prototype line clamp for venous access bleeding in hemodialysis: a prospective cohort study. *Hemodial Int.* 2019;23(2):151-157.
11. Agar JW, Perkins A, Heaf JG. Home hemodialysis: infrastructure, water, and machines in the home. *Hemodial Int.* 2015;19(suppl 1):S93-S111.
12. Hamilton TE, Centers for Medicare & Medicaid Services. Survey guidance for a new home hemodialysis water treatment device, the “NxStage PureFlow™ SL Water Purification System (PureFlow™)”. Published August 24, 2007.
13. Burton JO, Jefferies HJ, Selby NM, McIntyre CW. Hemodialysis-induced cardiac injury: determinants and associated outcomes. *Clin J Am Soc Nephrol.* 2009;4(5):914-920.
14. Movilli E, Gaggia P, Zubani R, et al. Association between high ultrafiltration rates and mortality in uraemic patients on regular haemodialysis. A 5-year prospective observational multicentre study. *Nephrol Dial Transplant.* 2007;22(12):3547-3552.
15. FHN Trial Group, Chertow GM, Levin NW, et al. In-center hemodialysis six times per week versus three times per week. *N Engl J Med.* 2010;363(24):2287-2300.
16. Rocco MV, Lockridge RS Jr, Beck GJ, et al. The effects of frequent nocturnal home hemodialysis: the Frequent Hemodialysis Network nocturnal trial. *Kidney Int.* 2011;80(10):1080-1091.

17. Jun M, Jardine MJ, Gray N, et al. Outcomes of extended-hours hemodialysis performed predominantly at home. *Am J Kidney Dis.* 2013;61(2):247-253.
18. Weinhandl ED, Nieman KM, Gilbertson DT, Collins AJ. Hospitalization in daily home hemodialysis and matched thrice-weekly in-center hemodialysis patients. *Am J Kidney Dis.* 2015;65(1):98-108.
19. Van Eps CL, Jones M, Ng T, et al. The impact of extended-hours home hemodialysis and buttonhole cannulation technique on hospitalization rates for septic events related to dialysis access. *Hemodial Int.* 2010;14(4):451-463.
20. Mustafa RA, Zimmerman D, Rioux JP, et al. Vascular access for intensive maintenance hemodialysis: a systematic review for a Canadian Society of Nephrology clinical practice guideline. *Am J Kidney Dis.* 2013;62(1):112-131.
21. MacRae JM, Ahmed SB, Atkar R, Hemmelgarn BR. A randomized trial comparing buttonhole with rope ladder needling in conventional hemodialysis patients. *Clin J Am Soc Nephrol.* 2012;7(10):1632-1638.
22. Nesrallah GE, Cuerden M, Wong JH, Pierratos A. *Staphylococcus aureus* bacteremia and buttonhole cannulation: long-term safety and efficacy of mupirocin prophylaxis. *Clin J Am Soc Nephrol.* 2010;5(6):1047-1053.
23. Daugirdas JT, Greene T, Rocco MV, et al. Effect of frequent hemodialysis on residual kidney function. *Kidney Int.* 2013;83(5):949-958.
24. Locatelli F, La Milia V, Violo L, Del Vecchio L, Di Filippo S. Optimizing haemodialysate composition. *Clin Kidney J.* 2015;8(5):580-589.
25. Kim SJ, Goldstein M, Szabo T, Pierratos A. Resolution of massive uremic tumoral calcinosis with daily nocturnal home hemodialysis. *Am J Kidney Dis.* 2003;41(3):E12.
26. Walsh M, Manns BJ, Klarenbach S, Tonelli M, Hemmelgarn B, Culleton B. The effects of nocturnal compared with conventional hemodialysis on mineral metabolism: a randomized-controlled trial. *Hemodial Int.* 2010;14(2):174-181.
27. Al-Hejaili F, Kortas C, Leitch R, et al. Nocturnal but not short hours quotidian hemodialysis requires an elevated dialysate calcium concentration. *J Am Soc Nephrol.* 2003;14(9):2322-2328.
28. Kayser GA, Greene T, Larive B, et al. The effect of frequent hemodialysis on nutrition and body composition: Frequent Hemodialysis Network trial. *Kidney 29.* Merlino G, Piani A, Dolso P, et al. Sleep disorders in patients with end-stage renal disease undergoing dialysis therapy. *Nephrol Dial Transplant.* 2006;21(1):184-190.
30. Jaber BL, Schiller B, Burkart JM, et al. Impact of short daily hemodialysis on restless legs symptoms and sleep disturbances. *Clin J Am Soc Nephrol.* 2011;6(5):1049-1056.
31. Pressman MR, Benz RL, Schleifer CR, Peterson DD. Sleep disordered breathing in ESRD: acute beneficial effects of treatment with nasal continuous positive airway pressure. *Kidney Int.* 1993;43(5):1134-1139.
32. Beecroft JM, Hoffstein V, Pierratos A, Chan CT, McFarlane P, Hanly PJ. Nocturnal haemodialysis increases pharyngeal size in patients with sleep apnoea and end-stage renal disease. *Nephrol Dial Transplant.* 2008;23(2):673-679.
33. Nearhos J, van Eps C, Connor J. Psychological factors associated with successful outcomes in home haemodialysis. *Nephrology (Carlton).* 2013;18(7):505-509.
34. Bowman B, Zheng S, Yang A, et al. Improving incident ESRD care via a transitional care unit. *Am J Kidney Dis.* 2018;72(2):278-283.
35. Suri RS, Larive B, Hall Y, et al. Effects of frequent hemodialysis on perceived caregiver burden in the Frequent Hemodialysis Network trials. *Clin J Am Soc Nephrol.* 2014;9(5):936-942.