

36.

BÖLÜM

BÖBREK VASKÜLER HASTALIKLARI VE BÖBREK YETMEZLİKLERİ HAYVAN MODELLERİ

Muhammed GÖMEÇ¹

Bu bölümde bir halk sağlığı problemi haline gelmiş olan böbrek yetmezliğini ilgilendiren renal arter ve veni de içeren hayvan modellerinden bahsedilecektir.

1- AKUT BÖBREK HASARI MODELLERİ

Akut böbrek yetmezliği (ABY), akut tübüler nekrozda denilen bir klinik duruma bağlı oluşan hızlı böbrek fonksiyon kaybı ile karakterizedir. ABY hastanede yatan hastaların % 10-20'sini etkiler ve bu hastaların da % 10'u renal replasman tedavisi gerektirir (1). Kritik hastalar arasında, akut böbrek hasarı (ABH) insidansı %45-50 gibi yüksek bir oranda bildirilmiştir (2). ABH, hastanede kalış süresini uzatarak maliyetlerde artışa neden olmaktadır. Kronik böbrek hastalığı (KBH) ve son dönem böbrek yetmezliğinde dâhil olmak üzere uzun yıllar tedivi gerektiren önemli bir mortalite ve morbidite nedenidi (3). Klinik hastalarda ABH'in başlangıcı ve ilerlemesinde iskemi, toksinler ve sepsis rol oynayabilir. Hospitalizasyonun giderek arttığı ve buna bağlı ABH gelişen hasta sayısının da artmaya devam ettiği göz önünde bulundurulduğunda bu hasarın sebeplerinin anlaşılması yukarıda说得larımız mortalite ve morbiditenin azalmasına hizmet edecektir. Bir çok alanda olduğu gibi bu konuyu anlamak için de hayvan modelleri ile çalışmalar yapılmaktadır. Genellikle bu üç etken iç içedir ve buna bağlı hayvan deney modellerini birebir klinikle özdeşleştirmek zordur. Klinik ABH'nın ana nedenlerine dayanarak, iskemi-reperfüzyon, toksinler ve sepsis olmak üzere üç tip hayvan modeli geliştirilmiştir.

¹ Dr. Öğr. Üyesi, Sivas Cumhuriyet Üniversitesi Tıp Fakültesi, Genel Cerrahi AD

KAYNAKLAR

1. Thongprayoon, C., Hansrivijit, P., Kovvuru, K., Kanduri, S. R., Torres-Ortiz, A., Acharya, P., . . . Cheungpasitporn, W. (2020). Diagnostics, Risk Factors, Treatment and Outcomes of Acute Kidney Injury in a New Paradigm. *J Clin Med*, 9(4). doi:10.3390/jcm9041104
2. Ronco, C., Bellomo, R., & Kellum, J. A. (2019). Acute kidney injury. *Lancet*, 394(10212), 1949-1964. doi:10.1016/s0140-6736(19)32563-2
3. Silver, S. A., Long, J., Zheng, Y., & Chertow, G. M. (2017). Cost of Acute Kidney Injury in Hospitalized Patients. *J Hosp Med*, 12(2), 70-76. doi:10.12788/jhm.2683
4. Martins, P. N., Chandraker, A., & Tullius, S. G. (2006). Modifying graft immunogenicity and immune response prior to transplantation: potential clinical applications of donor and graft treatment. *Transplant international*, 19(5), 351-359.
5. Balint, P., Chatel, R., Fekete, A., & Forgaçs, I. (1964). Haemodynamics and oxygen consumption of the kidney in Post-ischaemic renal failure. *Clin Sci*, 26, 471-477.
6. Shoskes, D. A., Parfrey, N. A., & Halloran, P. F. (1990). Increased major histocompatibility complex antigen expression in unilateral ischemic acute tubular necrosis in the mouse. *Transplantation*, 49(1), 201-207. doi:10.1097/00007890-199001000-00045
7. Chen, H., Xing, B., Liu, X., Zhan, B., Zhou, J., Zhu, H., & Chen, Z. (2008). Ozone oxidative preconditioning inhibits inflammation and apoptosis in a rat model of renal
8. Kocoglu, H., Ozturk, H., Ozturk, H., Yilmaz, F., & Gulcu, N. (2009). Effect of dexmedetomidine on ischemia-reperfusion injury in rat kidney: a histopathologic study. *Renal failure*, 31(1), 70-74.
9. Doi, K., Suzuki, Y., Nakao, A., Fujita, T., & Noiri, E. (2004). Radical scavenger edaravone developed for clinical use ameliorates ischemia/reperfusion injury in rat kidney. *Kidney international*, 65(5), 1714-1723.
10. Matsuyama, M., Hayama, T., Funao, K., Tsuchida, K., Takemoto, Y., Sugimura, K., . . . Yoshimura, R. (2006). Treatment with edaravone improves the survival rate in renal warm ischemia-reperfusion injury using rat model. Paper presented at the Transplantation proceedings.
11. Meldrum, K. K., Meldrum, D. R., Meng, X., Ao, L., & Harken, A. H. (2002). TNF- α -dependent bilateral renal injury is induced by unilateral renal ischemia-reperfusion. *American Journal of Physiology-Heart and Circulatory Physiology*, 282(2), H540-H546.
12. Nezamoleslami, S., Sheibani, M., Dehpour, A. R., Mobasheran, P., & Shafaroodi, H. (2020). Glatiramer acetate attenuates renal ischemia reperfusion injury in rat model. *Experimental and Molecular Pathology*, 112, 104329.
13. Abdallah, N. H., Baülies, A., Bouhlel, A., Bejaoui, M., Zaouali, M. A., Mimouna, S. B., . . . Abdennabi, H. B. (2018). The effect of zinc acexamate on oxidative stress, inflammation and mitochondria induced apoptosis in rat model of renal warm ischemia. *Biomedicine & Pharmacotherapy*, 105, 573-581.
14. Ju, C.-h., Xue, L.-n., & Cheng, H.-j. (2018). Optimized method to harvest both kidneys from one donor rat for transplantation. *BMC surgery*, 18(1), 1-6.
15. Harvig, B., Engberg, A., & Ericsson, J. L. (1980). Effects of cold ischemia on the preserved and transplanted rat kidney. Structural changes of the loop of Henle, distal tubule and collecting duct. *Virchows Arch B Cell Pathol Incl Mol Pathol*, 34(2), 173-192. doi:10.1007/bf02892416
16. Sayed-Ahmed, M. M., Eissa, M. A., Kenawy, S. A., Mostafa, N., Calvani, M., & Osman, A.-M. M. (2004). Progression of cisplatin-induced nephrotoxicity in a carnitine-depleted rat model. *Cancer chemotherapy*, 50(4), 162-170.
17. Wu, C. T., Sheu, M. L., Tsai, K. S., Chiang, C. K., & Liu, S. H. (2011). Salubrin, an eIF2 α dephosphorylation inhibitor, enhances cisplatin-induced oxidative stress and nephrotoxicity in a mouse model. *Free Radical Biology and Medicine*, 51(3), 671-680.

18. Wei, Q., Wang, M. H., & Dong, Z. (2005). Differential gender differences in ischemic and nephrotoxic acute renal failure. *Am J Nephrol*, 25(5), 491-499. doi:10.1159/000088171
19. Liu, Y., Tolbert, E. M., Lin, L., Thursby, M. A., Sun, A. M., Nakamura, T., & Dworkin, L. D. (1999). Up-regulation of hepatocyte growth factor receptor: an amplification and targeting mechanism for hepatocyte growth factor action in acute renal failure. *Kidney international*, 55(2), 442-453.
20. Wan, B., Hao, L., Qiu, Y., Sun, Z., Cao, Q., Zhang, Y., . . . Zhang, Y. (2006). Blocking tumor necrosis factor- α inhibits folic acid-induced acute renal failure. *Experimental and Molecular Pathology*, 81(3), 211-216. doi:<https://doi.org/10.1016/j.yexmp.2006.02.005>
21. Santos, S., Bosch, R. J., Ortega, A., Largo, R., Fernández-Agulló, T., Gazapo, R., . . . Esbrí, P. (2001). Up-regulation of parathyroid hormone-related protein in folic acid-induced acute renal failure. *Kidney international*, 60(3), 982-995. doi:<https://doi.org/10.1046/j.1523-1755.2001.060003982.x>
22. Zhao, B., Zhao, Q., Li, J., Xing, T., Wang, F., & Wang, N. (2015). Renalase protects against contrast-induced nephropathy in Sprague-Dawley rats. *PloS one*, 10(1), e0116583.
23. Sun, S., Zhang, T., Nie, P., Hu, L., Yu, Y., Cui, M., . . . He, B. (2014). A novel rat model of contrast-induced acute kidney injury. *International Journal of Cardiology*, 172(1), e48-e50.
24. Miyaji, T., Hu, X., Yuen, P. S. T., Muramatsu, Y., Iyer, S., Hewitt, S. M., & Star, R. A. (2003). Ethyl pyruvate decreases sepsis-induced acute renal failure and multiple organ damage in aged mice. *Kidney international*, 64(5), 1620-1631. doi:<https://doi.org/10.1046/j.1523-1755.2003.00268.x>
25. Holly, M. K., Dear, J. W., Hu, X., Schechter, A. N., Gladwin, M. T., Hewitt, S. M., . . . Star, R. A. (2006). Biomarker and drug-target discovery using proteomics in a new rat model of sepsis-induced acute renal failure. *Kidney international*, 70(3), 496-506. doi:<https://doi.org/10.1038/sj.ki.5001575>
26. Ma, N., Wang, S. Y., Sun, Y. J., Ren, J. H., & Guo, F. J. (2019). [Diagnostic value of contrast-enhanced ultrasound for accessory renal artery among patients suspected of renal artery stenosis]. *Zhonghua Yi Xue Za Zhi*, 99(11), 838-840. doi:10.3760/cma.j.issn.0376-2491.2019.11.008
27. van Twist, D. J. L., Houben, A. J. H. M., de Haan, M. W., de Leeuw, P. W., & Kroon, A. A. (2017). Pathophysiological differences between multifocal fibromuscular dysplasia and atherosclerotic renal artery stenosis. *Journal of hypertension*, 35(4), 845-852. doi:10.1097/hjh.0000000000001243
28. Truong, L. D., Farhood, A., Tasby, J., & Gillum, D. (1992). Experimental chronic renal ischemia: morphologic and immunologic studies. *Kidney international*, 41(6), 1676-1689.
29. Konopka, C. L., Jurach, A., & Wender, O. C. B. (2007). Experimental model for the study of chronic renal ischemia in rats: morphologic, histological and ultra-structural analysis. *Acta Cirúrgica Brasileira*, 22(1), 12-21.
30. Molitch, M. E., DeFronzo, R. A., Franz, M. J., & Keane, W. F. (2004). Nephropathy in diabetes. *Diabetes care*, 27, S79.
31. Wang, S., Li, Y., Zhao, J., Zhang, J., & Huang, Y. (2013). Mesenchymal Stem Cells Ameliorate Podocyte Injury and Proteinuria in a Type 1 Diabetic Nephropathy Rat Model. *Biology of Blood and Marrow Transplantation*, 19(4), 538-546. doi:<https://doi.org/10.1016/j.bbmt.2013.01.001>
32. Candido, R., Jandeleit-Dahm, K. A., Cao, Z., Nesteroff, S. P., Burns, W. C., Twigg, S. M., . . . Allen, T. J. (2002). Prevention of accelerated atherosclerosis by angiotensin-converting enzyme inhibition in diabetic apolipoprotein E-deficient mice. *Circulation*, 106(2), 246-253.
33. Tesch, G. H., & Allen, T. J. (2007). Rodent models of streptozotocin-induced diabetic nephropathy (Methods in Renal Research). *Nephrology*, 12(3), 261-266.
34. Wu, M.-J., Wen, M.-C., Chiu, Y.-T., Chiou, Y.-Y., Shu, K.-H., & Tang, M.-J. (2006). Rapamycin attenuates unilateral ureteral obstruction-induced renal fibrosis. *Kidney international*, 69(11), 2029-2036.

35. KIM, W. Y., Nam, S. A., Song, H. C., Ko, J. S., Park, S. H., Kim, H. L., . . . Kim, Y. K. (2012). The role of autophagy in unilateral ureteral obstruction rat model. *Nephrology*, 17(2), 148-159.
36. Xianyuan, L., Wei, Z., Yaqian, D., Dan, Z., Xueli, T., Zhanglu, D., . . . Menghua, L. (2019). Anti-renal fibrosis effect of asperulosidic acid via TGF- β 1/smad2/smad3 and NF- κ B signaling pathways in a rat model of unilateral ureteral obstruction. *Phytomedicine*, 53, 274-285.
37. Li, C., Yang, C. W., Park, J. H., Lim, S. W., Sun, B. K., Jung, J. Y., . . . Bang, B. K. (2004). Pravastatin treatment attenuates interstitial inflammation and fibrosis in a rat model of chronic cyclosporine-induced nephropathy. *American journal of physiology-renal physiology*, 286(1), F46-F57.
38. Khan, M., Shobha, J. C., Mohan, I. K., Rao Naidu, M. U., Prayag, A., & Kutala, V. K. (2006). Spirulina attenuates cyclosporine-induced nephrotoxicity in rats. *Journal of Applied Toxicology: An International Journal*, 26(5), 444-451.
39. Saxton, N. E., Barclay, J. L., Clouston, A. D., & Fawcett, J. (2002). Cyclosporin A pretreatment in a rat model of warm ischaemia/reperfusion injury. *Journal of hepatology*, 36(2), 241-247.
40. Young, B. A., Burdmann, E. A., Johnson, R. J., Andoh, T., Bennett, W. M., Couser, W. G., & Alpers, C. E. (1995). Cyclosporine A induced arteriolopathy in a rat model of chronic cyclosporine nephropathy. *Kidney international*, 48(2), 431-438.