

17. BÖLÜM

PSİKIYATRİK HASTALIKLarda PREKONSEPSİYONEL BAKİM VE DANIŞMANLIK

Hacer YALNIZ DİLÇEN ¹
Birsen KARACA SAYDAM ²

GİRİŞ

Psikiyatrik hastalıklarda prekonsepsiyonel bakım, gebelik planlayan çiftlerin, gebelikten önce, sağlığını geliştirmeyi amaçlayan koruyucu bir sağlık hizmeti türü olup, tarama ile tedavinin yanı sıra danışmanlık hizmetini de içerir. Amacı, psikiyatrik hastalıkların neden olacağı, anne ve fetüs için risk oluşturan tıbbi ve sosyal durumları gebelikten önce saptayarak uygun şekilde çözmek veya yönlendirmektir. Bu dönem planlanmayan gebeliğin önlenmesi için önemli bir fırsattır. Planlanan bir gebeliği ise en sağlıklı şekilde başlatabilmek yine prekonsepsiyonel bakımla mümkündür.

Üreme çağındaki kadınların önemli bir kısmı için gebelik öncesi, gebelik, doğum ve doğum sonrası dönemler, psikiyatrik rahatsızlıkların ortaya çıkması veya önceden var olan bir psikiyatrik rahatsızlığın yinelemesi nedeniyle komplike olabilir. Gebelik sırasında ortaya çıkan psikiyatrik bozukluklar, obstetrik sonuçların kötü olması, postpartum psikiyatrik hastalık riski yüksekliği, madde kötüye kullanım oranlarının artması, doğum öncesi bakıma katılım oranının düşük olması, bebek ve ailede istenmeyen olumsuz sonuçlarla ilişkilidir.

Türkiye ruh sağlığı profili çalışmasına göre; ülkemizde ruhsal hastalıkların bir yıllık yaygınlığı %17.2'dir. Ortalama her beş kadından biri, son bir yıl içinde bir psikiyatrik hastalık geçirmiştir. Bunun yanı sıra araştırmalar, her iki kişiden birinin yaşam boyunca en az bir psikiyatrik hastalık yaşadığına işaret etmektedir. 15-49 yaş üreme çağında veya gebelik öncesi kadınlar arasında psikiyatrik semptomların saptanması ve uygun şekilde yönetimi; gebelik sırasında ve sonrasında morbiditeyi önlemek açısından kritik bir önem taşır. Gebelik öncesi

¹ Dr. Öğr. Üyesi, Bartın Üniversitesi, Sağlık Bilimleri Fakültesi, Ebelik Bölümü

² Doç. Dr., Ege Üniversitesi, Sağlık Bilimleri Fakültesi, Ebelik Bölümü

KAYNAKLAR

- Alwan S., Reehuis J., Rasmussen S.A., Olney R.S., Friedman J.M. (2007). Use of selective serotonin-reuptake inhibitors in pregnancy and the risk of birth defects. *The New England Journal of Medicine*, 356(26):2684-2692. Doi: 10.1056/NEJMoa066584
- American Psychiatric Association. (2013). Diagnostic and Statistical Manual of Mental Disorders, 5th Edition. Doi: 10.1176/appi.books.9780890425596.744053
- Amrollahi Z., Rezaei F., Salehi B., Modabbernia A.H., Maroufi A., Esfandiari G.R., ... Akhondzadeh S. (2011). Double-blind, randomized, placebo-controlled 6-week study on the efficacy and safety of the tamoxifen adjunctive to lithium in acute bipolar mania. *Journal of Affective Disorders*, 129(1-3):327-331.
- Avni-Barron O., Hoagland K., Ford C., Miller L.J. (2010). Preconception planning to reduce the risk of perinatal depression and anxiety disorders. *Expert Review of Obstetrics and Gynecology*, 5(4):421-435. Available at: <https://doi.org/10.1586/eog.10.27>
- Binbay T., Ulaş H., Elbi H., Alptekin K. (2011). Türkiye'de psikoz epidemiyolojisi: yaygınlık tahminleri ve başvuru oranları üzerine sistematik bir gözden geçirme. *Türk Psikiyatri Dergisi*, 22(1):40-52.
- Boden R., Lundgren M., Brandt L., Reutfors J., Andersen M., Kieler H. (2012). Risks of adverse pregnancy and birth outcomes in women treated or not treated with mood stabilisers for bipolar disorder: population based cohort study. *BMJ*, 345:e7085. Available at: <https://doi.org/10.1136/bmj.e7085>
- Bonari L., Pinto N., Ahn E., Einarson A., Steiner M., Koren G. (2004). Perinatal risks of untreated depression during pregnancy. *Canadian Journal of Psychiatry*, 49(11):726-735. Available at: <https://doi.org/10.1111/j.1399-3046.2011.01554.x>
- Bonnin C.M., Sinchez-Moreno J., Martinez-Arin A., Soli B., Reinares M., Rosa A.R., ... Torrent C. (2012). Subthreshold symptoms in bipolar disorder: Impact on neurocognition, quality of life and disability. *Journal of Affective Disorders*, 136(3):650-659. Available at: <https://doi.org/10.1016/j.jad.2011.10.012>
- Cantilino A., Lorenzo L., de Paula J. dos A., Einarson A. (2014). Use of psychotropic medications during pregnancy: perception of teratogenic risk among physicians in two Latin American countries. *Revista Brasileira de Psiquiatria*, 36(2):106-110. Available at: <https://doi.org/10.1590/1516-4446-2013-1221>
- Cohen L.S., Wang B., Nonacs R., Viguera A.C., Lemon E.L., Freeman M.P. (2010). Treatment of mood disorders during pregnancy and postpartum. *Psychiatric Clinics of North America*, 33(2):273-293. Available at: <https://doi.org/10.1016/j.psc.2010.02.001>
- Çetin M. (2010). Sızofreni tedavisinde gelinen nokta: tipik ve yeni antipsikotikler ne kadar başlırlar? *Turkiye Klinikleri Journal of Psychiatry Special Topics*, 3(2):45-53.
- de Graaf R., Ten Have M., van Gool C., van Dorsselaer S. (2012). Prevalence of mental disorders, and trends from 1996 to 2009. *Tijdschrift Voor Psychiatrie*, 54(1):27-38.
- Dennis C.L., McQueen K. (2009). The relationship between infant-feeding outcomes and postpartum depression: a qualitative systematic review. *Pediatrics*, 123(4):e736-51. Available at: <https://doi.org/10.1542/peds.2008-1629>
- Depp C.A., Jeste D.V. (2004). Bipolar disorder in older adults: a critical review. *Bipolar Disorders*, 6(5):343-367. Available at: <https://doi.org/10.1111/j.1399-5618.2004.00139.x>
- Farahi N., Zolotor A. (2013). Recommendations for preconception counseling and care. *American Family Physician*, 88(8):499-506.
- Ferri C.P., Mitsuhiro S.S., Barros M.C., Chalem E., Guinsburg R., Patel V., ... Laranjeira R. (2007). The impact of maternal experience of violence and common mental disorders on neonatal outcomes: a survey of adolescent mothers in São Paulo, Brazil. *BMC Public Health*, 7(1):209. Available at: <https://doi.org/10.1186/1471-2458-7-209>
- Fornaro M., Orsolini L., Marini S., De Berardis D., Perna G., Valchera A., ... Stubbs B. (2016). The prevalence and predictors of bipolar and borderline personality disorders comorbidity:

- systematic review and meta-analysis. *Journal of Affective Disorders*, 195:105-118. Available at: <https://doi.org/10.1016/j.jad.2016.01.040>
- Frayne J., Nguyen T., Allen S., Rampono J. (2009). Motherhood and mental illness: part 1 - toward a general understanding. *Australian Family Physician*, 38(8):594-600.
- Freeman M.P. (2007). Bipolar disorder and pregnancy: risks revealed. *American Journal of Psychiatry*, 164(12):1771-1773. Doi: 10.1176/appi.ajp.2007.07091408
- Frieder A., Dunlop A.L., Culpepper L., Bernstein P.S. (2008). The clinical content of preconception care: women with psychiatric conditions. *American Journal of Obstetrics and Gynecology*, 199(6 SUPPL. B):328-332. Doi: 10.1016/j.ajog.2008.09.001
- Friedman S.H., Resnick P.J. (2009). Postpartum depression: an update. *Women's Health*, 5(3):287-295. Available at: <https://doi.org/10.2217/whe.09.3>
- Gavin N., Gaynes B. (2005). Perinatal depression: a systematic review of prevalence and incidence. *Journal of Obstetrics and Gynecology*, 106(5 Pt 1):1071-1083.
- Halter M.J. (2018). Varcarolis' Foundations of Psychiatric Mental Health Nursing : A Clinical Approach, 8th Edition, Elsevier: St. Louis.
- Hendrick V.C. (2017). Bipolar disorder in women: contraception and preconception assessment and counseling. Available at: <https://www.uptodate.com/contents/bipolar-disorder-in-women-contraception-and-preconception-assessment-and-counseling>
- Hirani S.S., Karmaliani R., McFarlane J., Asad N., Madhani F., Shehzad S. (2010). Testing a community derived intervention to promote women's health: preliminary results of a 3-arm randomized controlled trial in Karachi, Pakistan. *Southern Online Journal of Nursing Research*, 10(3):1-5.
- Howard L.M., Bekele D., Rowe M., Demilew J., Bewley S., Marteau T.M. (2013). Smoking cessation in pregnant women with mental disorders: a cohort and nested qualitative study. *International Journal of Obstetrics and Gynaecology*, 120(3):362-370. Available at: <https://doi.org/10.1111/1471-0528.12059>
- Howard L.M., Megnin-Viggars O., Symington I., Pilling S. (2014). Antenatal and postnatal mental health: summary of updated NICE guidance. *BMJ*, 349:g7394-g7394. Available at: <https://doi.org/10.1136/bmj.g7394>
- Huybrechts K.F., Hernández-Díaz S., Patorno E., Desai R.J., Mogun H., Dejene S.Z., ... Bateman B.T. (2016). Antipsychotic use in pregnancy and the risk for congenital malformations. *JAMA Psychiatry*, 66(4):1-9.
- Johnson K., Posner S.F., Biermann M.S., Cordero J.F., Atrash H.K., Parker C.S., ... Curtis, M.S. (2006). Recommendations to improve preconception health and health care: strategies for implementation. A Report of the CDC/ATSDR Preconception Care Work Group and the Select Panel on Preconception Care, CDC MMWR, 55(RR-6). Available at: <https://doi.org/10.1089/jwh.2007.CDC3>
- Jones H.E., Kaltenbach K., Heil S.H., Stine S.M., Coyle M.G., Arria A.M., ... Fischer G. (2010). Neonatal abstinence syndrome after methadone or buprenorphine exposure. *The New England Journal of Medicine*, 363(24):2320-2331. Available at: <https://doi.org/10.1056/NEJMoa1005359>
- Sadock J.B., Sadock A.V., Ruiz P. (2016). Kaplan & Sadock's Comprehensive Textbook of Psychiatry, 11th Ed., Çev. Ed. Bozkurt A., Güneş Tip Kitap Evleri, ss.300-413.
- Karaçam Z., Ançel G. (2009). Depression, anxiety and influencing factors in pregnancy: a study in a Turkish population. *Midwifery*, 25(4):344-356.
- King-Hele S., Webb R.T., Mortensen P.B., Appleby L., Pickles A., Abel K.M. (2009). Risk of stillbirth and neonatal death linked with maternal mental illness: a national cohort study. *Archives of Disease in Childhood Fetal and Neonatal Edition*, 94(2):F105-110. Available at: <https://doi.org/10.1136/adc.2007.135459>
- Lassi Z.S., Imam A.M., Dean S.V., Bhutta Z.A. (2014). Preconception care: caffeine, smoking, alcohol, drugs and other environmental chemical/radiation exposure. *Reproductive Health*, 11(Suppl 3): S6. Doi: 10.1186/1742-4755-11-S3-S6

- Lin H.C., Chen I.J., Chen Y.H., Lee H.C., Wu F.J. (2010). Maternal schizophrenia and pregnancy outcome: does the use of antipsychotics make a difference? *Schizophrenia Research*, 116(1):55-60. Doi: 10.1016/j.schres.2009.10.011
- McAllister-Williams R.H., Baldwin D.S., Cantwell R., Easter A., Gilvarry E., Glover V., ... Young A.H. (2017). British Association for Psychopharmacology consensus guidance on the use of psychotropic medication preconception, in pregnancy and postpartum 2017. *Journal of Psychopharmacology*, 31(5):519-552.
- Meltzer-Brody S., Jones I. (2015). Optimizing the treatment of mood disorders in the perinatal period. *Dialogues in Clinical Neuroscience*, 17(2):207-218.
- Mennes M., den Bergh V.B., Lagae L., Stiers P. (2009). Developmental brain alterations in 17 year old boys are related to antenatal maternal anxiety. *Clinical Neurophysiology*, 120(6):1116-1122. Doi: 10.1016/j.clinph.2009.04.003
- Merikangas K.R., Jin R., He J.P., Kessler R.C., Lee S., Sampson N.A., ... Zarkov Z. (2011). Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative. *Archives of General Psychiatry*, 68(3):241-251.
- Moos M.K., Dunlop A.L., Jack B.W., Nelson L., Coonrod D.V., Long R., ... Gardiner P.M. (2008). Healthier women, healthier reproductive outcomes: recommendations for the routine care of all women of reproductive age. *American Journal of Obstetrics and Gynecology*, 199(6 Suppl 2):S280-S289. Doi: 10.1016/j.ajog.2008.08.060
- Munk-Olsen T., Laursen T.M., Mendelson T., Pedersen C.B., Mors O., Mortensen P.B. (2009). Risks and predictors of readmission for a mental disorder during the postpartum period. *Archives of General Psychiatry*, 66(2):189-195.
- Muzik M., Marcus S.M., Heringhausen J.E., Flynn H. (2009). When depression complicates childbearing: guidelines for screening and treatment during antenatal and postpartum obstetric care. *Obstetrics and Gynecology Clinics of North America*, 36(4):771-788. Available at: <https://doi.org/10.1016/j.ocg.2009.10.006>
- Parial S. (2015). Bipolar disorder in women. *Indian Journal of Psychiatry*, 57(Suppl 2):S252-263. <https://doi.org/10.4103/0019-5545.161488>
- Patton S.W., Misri S., Corral M.R., Perry K.F., Kan A.J. (2002). Antipsychotic medication during pregnancy and lactation in women with schizophrenia: evaluating the risk. *Canadian Journal of Psychiatry*, 47(10):959-965.
- RaiKKanen K., Lahti M., Heinonen K., Pesonen A. K., Wahlbeck K., Kajantie E., ... Eriksson J.G. (2011). Risk of severe mental disorders in adults separated temporarily from their parents in childhood: the Helsinki birth cohort study. *Journal of Psychiatric Research*, 45(3):332-338. Doi: 10.1016/j.jpsychires.2010.07.003
- Rychtarik R.G., McGillicuddy N.B., Barrick C. (2015). Web-based coping skills training for women whose partner has a drinking problem. *Psychology of Addictive Behaviors*, 29(1):26-33. Available at: <https://doi.org/10.1037/adb0000032>
- Sax H., Allegranzi B., Chraiti M.N., Boyce J., Larson E., Pittet D. (2009). The World Health Organization hand hygiene observation method. *American Journal of Infection Control*, 37(10):827-834. Available at: <https://doi.org/10.1016/j.ajic.2009.07.003>
- Schluter P.J., Lacey C., Porter R.J., Jamieson H.A. (2017). An epidemiological profile of bipolar disorder among older adults with complex needs: a national cross-sectional study. *Bipolar Disorders*, 19(5):375-385. Doi: 10.1111/bdi.12511
- Silverman M.E., Loudon H. (2010). Antenatal reports of pre-pregnancy abuse is associated with symptoms of depression in the postpartum period. *Archives of Women's Mental Health*, 13(5):411-415. Doi: 10.1007/s00737-010-0161-7
- Stevens A.W.M.M., Daggenvoorde T.H., van der Klis S.M.D., Kupka R.W., Goossens P.J.J. (2018). Thoughts and considerations of women with bipolar disorder about family planning and pregnancy: a qualitative study. *Journal of the American Psychiatric Nurses Association*, 24(2):118-126. Doi: 10.1177/1078390317711251

- The Royal College of Midwives. (2015). Caring for Women with Mental Health Problems, England, pp.1-32.
- Vega P., Barbeito S., Ruiz de Azúa S., Martínez-Cengotitabengoa M., González-Ortega I., Saenz M., González-Pinto A. (2011). Bipolar disorder differences between genders: special considerations for women. *Women's Health*, 7(6):663-674. Doi: 10.2217/whe.11.71
- Ververs T., van Dijk L., Yousofi S., Schobben F., Visser G.H.A. (2009). Depression during pregnancy: views on antidepressant use and information sources of general practitioners and pharmacists. *BMC Health Services Research*, 9:119. Doi: 10.1186/1472-6963-9-119
- Vigod S.N., Kurdyak P.A., Dennis C.L., Gruneir A., Newman A., Seeman M.V., ... Ray J.G. (2014). Maternal and newborn outcomes among women with schizophrenia: a retrospective population-based cohort study. *BJOG: An International Journal of Obstetrics and Gynaecology*, 121(5):566-574. Doi: 10.1111/1471-0528.12567
- Viguera A.C., Cohen L.S., Baldessarini R.J., Nonacs R. (2002a). Managing bipolar disorder during pregnancy: weighing the risks and benefits. *Canadian Journal of Psychiatry*, 47(5):426-436. Doi: 10.1007/s11245-004-5371-z
- Viguera A.C., Cohen L.S., Bouffard S., Hatch Whitfield T., Baldessarini R.J. (2002b). Reproductive decisions by women with bipolar disorder after prepregnancy psychiatric consultation. *American Journal of Psychiatry*, 159(12):2102-2104.
- Viguera A.C., Koukopoulos A., Muzina D.J., Baldessarini R.J. (2007). Teratogenicity and anti-convulsants: lessons from neurology to psychiatry. *Journal of Clinical Psychiatry*, 68:29-33.
- Viguera A.C., Tondo L., Koukopoulos A.E., Reginaldi D., Lepri B., Baldessarini R.J. (2011). Episodes of mood disorders in 2,252 pregnancies and postpartum periods. *American Journal of Psychiatry*, 168(11):1179-1185.
- Walton G.D., Ross L.E., Stewart D.E., Grigoriadis S., Dennis C.L., Vigod S. (2014). Decisional conflict among women considering antidepressant medication use in pregnancy. *Archives of Women's Mental Health*, 17(6):493-501. Doi: 10.1007/s00737-014-0448-1
- Webb R.T., Wicks S., Dalman C., Pickles A.R., Appleby L., Mortensen P.B., ... Abel K.M. (2010). Influence of environmental factors in higher risk of sudden infant death syndrome linked with parental mental illness. *Archives of General Psychiatry*, 67(1):69-77. Available at: <https://doi.org/10.1001/archgenpsychiatry.2009.172>
- Wesseloo R., Kamperman A.M., Munk-Olsen T., Pop V.J.M., Kushner S.A., Bergink V. (2016). Risk of postpartum relapse in bipolar disorder and postpartum psychosis: a systematic review and meta-analysis. *American Journal of Psychiatry*, 173(2):117-127. Available at: <https://doi.org/10.1176/appi.ajp.2015.15010124>
- Whiteford H.A., Degenhardt L., Rehm J., Baxter A.J., Ferrari A.J., Erskine H.E., ... Vos T. (2013). Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. *The Lancet*, 382(9904):1575-1586. Doi: 10.1016/S0140-6736(13)61611-6
- Wichman C.L. (2016). Managing your own mood lability: use of mood stabilizers and antipsychotics in pregnancy. *Current Psychiatry Reports*, 18(1):1.
- Wieck A., Kopparthi S., Sundaresh S., Wittkowski A. (2010). One-year outcome after preconception consultation in women with bipolar disorder. *Journal of Clinical Psychiatry*, 71(6):806. Available at: <https://doi.org/10.4088/JCP.09l0559yel>
- Yalnız H., Türkmen H., Karaca Saydam B., et. al. (2016). Gebelik ve psikiyatrik hastalıklar. Uluslararası Hakemli Kadın Hastalıkları ve Anne Çocuk Sağlığı Dergisi, 7:1-20. Available at: <https://doi.org/10.17367/JACSD.2016721957>
- Yonkers K.A., Wisner K.L., Stowe Z., Leibenluft E., Cohen L., Miller L., ... Altshuler L. (2004). Management of bipolar disorder during pregnancy and the postpartum period. *The American Journal of Psychiatry*, 161(4):608-620.
- Zhang L., Cao X.L., Wang B.S., Zheng W., Ungvari G.S., Ng C.H., ... Xiang Y.T. (2017). The prevalence of bipolar disorder in China: a meta-analysis. *Journal of Affective Disorders*, 207:413-421. Available at: <https://doi.org/10.1016/j.jad.2016.08.062>