



KAYNAKLAR

1. Ortiz-Neira, CL., Paolucci, EO., Donnon T. A meta-analysis of common risk factors associated with the diagnosis of developmental dysplasia of the hip in newborns. *European Journal of Radiology*; 2012;81(3):e344–e351.
2. Yamamuro, T., Ishida, K. Recent advances in the prevention, early diagnosis, and treatment of congenital dislocation of the hip in Japan. *Clinical Orthopaedics and Related Research*; 1984;(184):34–40.
3. Harris, WH. Etiology of osteoarthritis of the hip. *Clinical Orthopaedics and Related Research*; 1986;(213):20–33.
4. Corbett D. Information needs of parents of a child in a Pavlik harness. *Orthopaedic Nursing*; 1988, 8(1), 20–23.
5. Weinstein, SL. (2001). Developmental hip dysplasia and dislocation. In R. T. Morrissy & S. L. Weinstein (Eds.), *Lovell and Winter's pediatric orthopaedics* (pp. 905–956). Philadelphia: Lippincott Williams and Wilkins.
6. Sparks, L., Ortman, MR., Aubuchon, BS. Meeting the developmental needs of a child in a body cast. *Journal of Orthopaedic Nursing*; 2005, 9(1), 35–38.
7. Georgiadis, AG., Seeley, MA., Yellin, JL., et al. The presentation of Legg-CalvéPerthes disease in females. *The Journal of Children's Orthopaedics*; 2015;9:243–7.
8. Perry, DC., Bruce, CE., Pope, D., et al. Comorbidities in Perthes' disease: a case control study using the General Practice Research database. *Journal of bone and joint surgery: British*; 2012; 94:1684-9.
9. Cooperman, DR., Stulberg, SD. Ambulatory containment treatment in Perthes' disease. *Clinical Orthopaedics and Related Research*; 1986 Feb;203:289-300.
10. Meehan, PL., Angel, D., Nelson, JM. The Scottish Rite abduction orthosis for the treatment of Legg-Perthes disease. A radiographic analysis. *Journal of bone and joint surgery: American*; 1992 Jan;74(1):2-12.
11. Joseph, B., Nair, NS., Narasimha Rao, K., et al. Optimal timing for containment surgery for Perthes disease. *Journal of Pediatric Orthopaedics*; 2003 Sep-Oct;23(5):601-6.
12. Choi, IH., Shin, YW., Chung, CY., et al. Surgical treatment of the severe sequelae of infantile septic arthritis of the hip. *Clinical Orthopaedics and Related Research; Clin Orthop Relat Res*, 2005; (434): 102–9
13. Abdelaziz, TH., El-Sayed, MM. Pauwels' osteotomy for surgical correction of infantile coxa vara. *Journal of Pediatric Orthopaedics B*; 2012;21:325–330.
14. Sabharwal, S., Mittal, R., Cox, G. Percutaneous triplanar femoral osteotomy correction for developmental coxa vara: a new technique. *Journal of Pediatric Orthopaedics*; 2005; 25(1): 28-33.
15. Herring, JA. (2014). Congenital Coxa Vara. In: Herring JA, editor. *Tachdjian Pediatric Orthopedics*, 5th ed. Philadelphia: Elsevier; p.666–77.
16. Paley, D. (2005). *Principles of deformity correctin*. Chapter 20, 3rd ed. New York: SpringerVerlag Berlin Heidelberg, p.705.
17. Danino, B., Rodl, R., Herzenberg, JE., et al. Growth modulation in idiopathic angular knee deformities: is it predictable? *The Journal of Children's Orthopaedics*; 2019; 13: 318–323.
18. Canale, ST. (2003). *Campbell's operative orthopedics*, Vol. 3. 10th ed. Mosby, Philadelphia, 2165-323.
19. Insall, JN., Joseph, DM., Msika, C. High tibial osteotomy for varus gonarthrosis. A long-term follow-up study. *The Journal of bone and joint surgery. American*; 1984; 66: 1040-8.



20. Salenius, P., Vankka, E. The development of the tibiofemoral angle in children, *The Journal of bone and joint surgery. American*; 57 (2) (1975) 259–261.
21. Arazi, M., Ogün, TC., Memik, R. Normal development of the tibiofemoral angle in children: a clinical study of 590 normal subjects from 3 to 17 years of age, *Journal of Pediatric Orthopaedics*; 21 (2), 264–267.
22. Ganesan, B., Fong, KN., Luximon, A., et al. Kinetic and kinematic analysis of gait pattern of 13 year old children with unilateral genu valgum, *European Review for Medical and Pharmacological Sciences*; 2016, 20 (15), 3168–3171.
23. Andrews, M., Noyes, FR., Hewett, TE., et al. Lower limb alignment and foot angle are related to stance phase knee adduction in normal subjects: a critical analysis of the reliability of gait analysis data, *The Journal of Orthopaedic Research*; 1996, 14 (2) 289–295.
24. Van Gheluwe, B., Kirby, KA., Hagman, F. Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait. *Journal of the American Podiatric Medical Association*; 2005, 95 (6), 531–541.
25. Stevens, PM., MacWilliams, B., Mohr, RA. Gait analysis of stapling for genu valgum. *Journal of Pediatric Orthopaedics*; 2004, 24 (1), 70–74.
26. Pretkiewicz-Abacjew, E. Knock knee and the gait of six-year-old children. *The Journal of sports medicine and physical fitness*; 2005, 43 (2), 156–164.
27. Kearney, SP., Mosca, VS. Selective hemiepiphysiodesis for patellar instability with associated genu valgum. *Journal Orthopaedic*; 2015, 12(1):17–22.
28. King, S., Chambers, CT., Huguet, A., et al. The epidemiology of chronic pain in children and adolescents revisited: A systematic review. *Pain*. 2011; 152(12): 2729- 2738.
29. Lohrer, H., Nauck, T., Scholl, J., et al. Extracorporeal shock wave therapy for patients suffering from recalcitrant Osgood- Schlatter disease. *Sportverletzung – Sportschaden*; 2012, 26(4), 218e222.
30. Circi, E., Atalay, Y., Beyzadeoglu, T. Treatment of Osgood-Schlatter disease: review of the literature. *Musculoskelet Surg*. 2017; 101(3): 195- 200.
31. Gholve, PA., Scher, DM., Khakharia, S., et al. Osgood schlatter syndrome. *Current Opinion in Pediatrics*; 2007, 19(1), 44e50.
32. Nakase, J., Goshima, K., Numata, H., et al. Precise risk factors for Osgood-Schlatter disease. *Arch Orthop Trauma Surg*. 2015; 135(9): 1277- 1281.
33. Hirano, A., Fukubayashi ,T., Ishii, T., et al. Magnetic resonance imaging of Osgood-Schlatter disease: the course of the disease. *Skeletal Radiology*; 2002; 31(6): 334- 342.
34. Hefti, F. (2015). *Pediatric Orthopaedics in practice*. Springer.
35. Smith, JM., Varacallo, M. (2019). *Osgood Schlatter’s disease (tibial tubercle apophysitis)*. Stat-Pearls. Treasure Island
36. Topol, G., Podesta, L., Reeves, K., et al. Hyperosmolar dextrose injection for recalcitrant Osgood-Schlatter disease. *Pediatrics [Internet]*; 2011, 128(5), e1121ee1128.
37. Ošťádal, M., Chomiak, J., Dungal, P., et al. Comparison of the short-term and long-term results of the Ponseti method in the treatment of idiopathic pes equinovarus. *International Orthopaedics*, 2013; 37: 1821-1825.
38. Diméglio, A., Bonnet, F., Mazeau, P., et al. Ortopaedic treatment and passive motion machine: consequences for the surgical treatment of clubfoot. *Journal of Pediatric Orthopaedics*, 1996; 3: 173-180.



39. Dare, D., Dodwell, ER. Pediatric flatfoot: Cause and epidemiology, assessment and treatment. *Current Opinion in Pediatrics*; 2014;26(1):93–100.
40. Halabchi, F, Mazaheri, R., Mirshahi, M., et al. Pediatric flexible flatfoot; clinical aspects and algorithmic approach. *Iranian Journal of Pediatrics*; 2013;23(3):247–60.
41. MacKenzie, AJ, Rome, K., Evans, A. The efficacy of non-surgical interventions for pediatric flexible flat foot: A Critical Review. *Journal of Pediatric Orthopaedics*; 2012;32(8):830–34.
42. Ford, SE., Scannell, BP. Pediatric flatfoot: Pearls and pitfalls. *Foot and Ankle Clinics*; 2017;22(3):643–56.
43. Rome, K., Ashford, RL., Evans, A. Non-surgical interventions for paediatric pes planus. *Cochrane Database of Systematic Reviews*; 2010;(7):CD006311.
44. Turner, C., Gardiner, MD., Midgley, A., et al. A guide to the management of paediatric pes planus. *The Australian Journal of General Practice*; 2020 May;49(5):245-249.
45. Younger, AS., Hansen, ST. Adult cavovarus foot. *The Journal of the American Academy of Orthopaedic Surgeons*; 2005 Sep;13(5):302-15.
46. Manoli, A., Graham, B. The subtle cavus foot, “the underpronator”. *Foot and Ankle International*; 2005 Mar;26(3):256-63.
47. Ippolito, E. Update on pathologic anatomy of clubfoot. *The Journal of Pediatric Orthopaedics B*;1995;4(1):17-24.
48. Maynou, C., Szymanski, C., Thiounn, A. The adult cavus foot. *EFORT Open Reviews*; 2017 May;2(5):221-229.
49. Rosenbaum, AJ., Lisella, J., Patel, N., et al. The cavus foot. *Medical Clinics of North America*; 2014 Mar;98(2):301-12.
50. Burns, J., Raymond, J., Ouvrier, R. Feasibility of foot and ankle strength training in childhood Charcot-Marie-Tooth disease. *Neuromuscul Disorders*; 2009 Dec;19(12):818-21.
51. LoPiccolo, M., Chilvers, M., Graham, B., et al. Effectiveness of the cavus foot orthosis. *The Journal of Surgical Orthopaedic Advances*; 2010 Fall;19(3):166-9.
52. Krähenbühl, N., Weinberg, MW. Anatomy and Biomechanics of Cavovarus Deformity. *Foot and Ankle Clinics*; 2019 Jun;24(2):173-181.
53. Dobbs MB, Morcuendo JA. Other conditions of the hip. In: Lowel WW, Weinstein SL, Flynn JM, editors. *Lovell and Winter’s Pediatric Orthopaedics 7th ed.* Philadelphia: Wolters Kluwer Health/Lippincott Williams and Wilkins; 2013. p.1222–60.
54. Mayer NH, Esquenazi A, Childers MK. Common patterns of clinical motor dysfunction. *Muscle Nerve*; 1997; 6:(Suppl.) S21–S35.
55. Carr JB 2nd, Yang S, Lather LA. Pediatric pes planus: A state-of-the-art review. *Pediatrics* 2016;137(3):e20151230.
56. Stellwagen LM, Hubbard E, Vaux K. Look for the “stuck baby” to identify congenital torticollis. *Contemp Pediatr.* 2004;21(5):55–65
57. Ding J, Zhu T, Jin FC, et al. The effect of temporary hemiepiphysiodesis in the treatment of skeleton immature posttraumatic genu angular deformity: a retrospective study of 27 cases. *J Orthop Surg Res* 2019; 14: 381.