

DERİN VEN TROMBOZUNDA KOMPRESYON TEDAVİSİ

Barış AKÇA¹

Giriş

Pulmoner emboli (PE) ile birlikte venöz tromboembolizm klinik tablosunu oluşturan derin ven trombozu (DVT), alt ekstremitte veya üst ekstremitte derin venöz sistem elemanlarında ya da pelvik venlerde trombüs oluşması durumudur. Yapılan çalışmalarda, DVT insidansı yılda 70-140 vaka / 100.000 kişi olarak bildirilmiştir ⁽¹⁾.

Genel olarak, erken ve orta dönem komplikasyonları arasında trombozun ilerlemesi, PE ve DVT nüksü, uzun dönem komplikasyonları arasında, kronik venöz semptomlar ve / veya DVT'ye ikincil belirtiler olarak tanımlanan post-trombotik sendrom (PTS) sayılabilir. PTS Proksimal DVT'den sonraki 2 yıl içinde hastaların % 30-50'sinde ortaya çıkabilen en sık kronik DVT komplikasyonu olup (2). Olguların % 5-10'unda PTS şiddetlidir ⁽²⁾.

¹ Doç. Dr., İnönü Üniversitesi Tıp Fakültesi Turgut Özal Tıp Merkezi Kalp Damar Cerrahisi Anabilimdalı, dr.baris.akca@yandex.com.tr

Kaynaklar

1. Raskob GE, Angchaisuksiri P, Blanco AN, Buller H, Gallus A, Hunt BJ, Hylek EM, Kakkar A, Konstantinides SV, McCumber M, Ozaki Y, Wendelboe A, Weitz JI; ISTH Steering Committee for World Thrombosis Day. Thrombosis: a major contributor to global disease burden. *Arterioscler Thromb Vasc Biol* 2014;34:2363–71.
2. Prandoni P, Kahn SR. Post-thrombotic syndrome: prevalence, prognostication and need for progress. *Br J Haematol* 2009;145:286–295.
3. Korschake W, Valesky E, Stege H, Jünger M. Evidence of compression therapy. *Hautarzt*. 2017;68:625-31.
4. Altintas AA, Gehl B, Aust MC, Meyer-Marcotty M, Altintas MA. Impact of compression therapy on local microcirculation and histomorphology in venous leg ulcers. *Phlebologie*. 2011;40:9-14.
5. Beidler SK, Douillet CD, Berndt DF, Keagy BA, Rich PB, Marston WA. Inflammatory cytokine levels in chronic venous insufficiency ulcer tissue before and after compression therapy. *J Vasc Surg*. 2009;49:1013-20.
6. Beidler SK, Douillet CD, Berndt DF, Keagy BA, Rich PB, Marston WA. Multiplexed analysis of matrix metalloproteinases in leg ulcer tissue of patients with chronic venous insufficiency before and after compression therapy. *Wound Repair Regen*. 2008;16:642-8.
7. Dissemond J, Assenheimer B, Bültemann A, Gerber V, Gretener S, Kohler-von Siebenthal E, et al. Compression therapy in patients with venous leg ulcers. *J Dtsch Dermatol Ges*. 2016;14:1072-87.
8. Kahn SR, Shapiro S, Ducruet T, Wells PS, Rodger MA, Kovacs MJ, et al. Graduated compression stockings to treat acute leg pain associated with proximal DVT. A randomised controlled trial. *Thromb Haemost* 2014;112:1137-41.
9. Prandoni P, Lensing AW, Prins MH, Frulla M, Marchiori A, Bernardi E, et al. Below-knee elastic compression stockings to prevent the post-thrombotic syndrome: a randomized, controlled trial. *Ann Intern Med* 2004;141:249-56.
10. Ten Cate-Hoek AJ, Amin EE, Bouman AC, Meijer K, Tick LW, Middeldorp S, et al. Individualised versus standard duration of elastic compression therapy for prevention of post-thrombotic syndrome (IDEAL DVT): a multicentre, randomised, singleblind, allocation-concealed, non-inferiority trial. *Lancet Haematol* 2018;5:e25-33.
11. Partsch H, Kaulich M, Mayer W. Immediate mobilisation in acute vein thrombosis reduces post-thrombotic syndrome. *Int Angiol* 2004;23:206-12.
12. Prandoni P, Noventa F, Quintavalla R, Bova C, Cosmi B, Siragusa S, et al. Thigh-length versus below-knee compression elastic stockings for prevention of the postthrombotic syndrome in patients with proximal-venous thrombosis: a randomized trial. *Blood* 2012;119:1561-5.
13. Gloviczki P, Comerota AJ, Dalsing MC, Eklof BG, Gillespie DL, Gloviczki ML, et al. The care of patients with varicose veins and associated chronic venous diseases: clinical practice guidelines of the Society for Vascular Surgery and the American Venous Forum. *J Vasc Surg* 2011;53:2S-48S.
14. Mosti G, Iabichella ML, Partsch H. Compression therapy in mixed ulcers increases venous output and arterial perfusion. *J Vasc Surg* 2012;55:122-8.
15. Flour M, Clark M, Partsch H, Mosti G, Uhl JF, Chauveau M, et al. Dogmas and controversies in compression therapy: report of an International Compression Club (ICC) meeting, Brussels, May 2011. *Int Wound J* 2013;10:516-26.
16. Partsch H, Blattler W. Compression and walking versus bed rest in the treatment of proximal deep venous thrombosis with low molecular weight heparin. *J Vasc Surg* 2000;32:861-9.

17. Roumen-Klappe EM, den Heijer M, van Rossum J, Wollersheim H, van der Vleuten C, Thien T, et al. Multilayer compression bandaging in the acute phase of deep-vein thrombosis has no effect on the development of the post-thrombotic syndrome. *J Thromb Thrombolysis* 2009;27:400-5.
18. Arpaia G, Cimminiello C, Mastrogiacomo O, de Gaudenzi E. Efficacy of elastic compression stockings used early or after resolution of the edema on recanalization after deep venous thrombosis: the COMPRES Trial. *Blood Coagul Fibrinolysis* 2007;18:131-7.
19. Amin EE, Bistervels IM, Meijer K, Tick LW, Middeldorp S, Mostard G, et al. Reduced incidence of vein occlusion and postthrombotic syndrome after immediate compression for deep vein thrombosis. *Blood* 2018;132:2298-304.
20. Amin EE, Joore MA, Ten Cate H, Meijer K, Tick LW, Middeldorp S, et al. Clinical and economic impact of compression in the acute phase of deep vein thrombosis. *J Thromb Haemost* 2018;16:1555-63.
21. Donadini MP, Ageno W, Antonucci E, Cosmi B, Kovacs MJ, Le Gal G, et al. Prognostic significance of residual venous obstruction in patients with treated unprovoked deep vein thrombosis: a patient-level meta-analysis. *Thromb Haemost* 2014;111:172-9.
22. Kahn SR, Hirsch A, Shrier I. Effect of postthrombotic syndrome on health-related quality of life after deep venous thrombosis. *Arch Intern Med* 2002;162:1144-8.
23. Bergqvist D, Jendteg S, Johansen L, Persson U, Odegaard K. Cost of long-term complications of deep venous thrombosis of the lower extremities: an analysis of a defined patient population in Sweden. *Ann Intern Med* 1997;126:454-7.
24. Lubberts B, Paulino Pereira NR, Kabrhel C, Kuter DJ, DiGiovanni CW. What is the effect of venous thromboembolism and related complications on patient reported health-related quality of life? A meta-analysis. *Thromb Haemost* 2016;116:417-31.
25. Prandoni P, Lensing AW, Prins MH, Pesavento R, Piccioli A, Sartori MT, et al. The impact of residual thrombosis on the long-term outcome of patients with deep venous thrombosis treated with conventional anticoagulation. *Semin Thromb Hemost* 2015;41:133e40.
26. Strandness Jr DE, Langlois Y, Cramer M, Randlett A, Thiele BL. Long-term sequelae of acute venous thrombosis. *JAMA* 1983;250:1289-92.
27. Brandjes DP, Buller HR, Heijboer H, Huisman MV, de Rijk M, Jagt H, et al. Randomised trial of effect of compression stockings in patients with symptomatic proximal-vein thrombosis. *Lancet* 1997;349:759-62.
28. Kahn SR, Shapiro S, Wells PS, Rodger MA, Kovacs MJ, Anderson DR, et al. Compression stockings to prevent postthrombotic syndrome: a randomised placebo-controlled trial. *Lancet* 2014;383:880-8.
29. Jayaraj A, Meissner M. Impact of graduated compression stockings on the prevention of post-thrombotic syndrome-results of a randomized controlled trial. *Phlebology* 2015;30:541-8.
30. Appelen D, van Loo E, Prins MH, Neumann MH, Kolbach DN. Compression therapy for prevention of post-thrombotic syndrome. *Cochrane Database Syst Rev* 2017;9:CD004174.
31. Subbiah R, Aggarwal V, Zhao H, Kolluri R, Chatterjee S, Bashir R. Effect of compression stockings on post thrombotic syndrome in patients with deep vein thrombosis: a meta-analysis of randomised controlled trials. *Lancet Haematol* 2016;3:e293-300.
32. Kakkos SK, Daskalopoulou SS, Daskalopoulos ME, Nicolaidis AN, Geroulakos G. Review on the value of graduated elastic compression stockings after deep vein thrombosis. *Thromb Haemost* 2006;96:441-5.
33. Aschwanden M, Jeanneret C, Koller MT, Thalhammer C, Bucher HC, Jaeger KA. Effect of prolonged treatment with compression stockings to prevent post-thrombotic sequelae: a randomized controlled trial. *J Vasc Surg* 2008;47:1015- 21.

34. Mol GC, van de Ree MA, Klok FA, Tegelberg MJ, Sanders FB, Koppen S, et al. One versus two years of elastic compression stockings for prevention of post-thrombotic syndrome (OCTAVIA study): randomised controlled trial. *BMJ* 2016;353:i2691.
35. Ten Cate-Hoek AJ, Bouman AC, Joore MA, Prins M, Ten Cate H. for the IDEAL DVT Trial investigators. The IDEAL DVT study, individualised duration elastic compression therapy against long-term duration of therapy for the prevention of postthrombotic syndrome: protocol of a randomised controlled trial. *BMJ Open* 2014;4:e005265.
36. Kakkos SK, Gohel M, Baekgaard N, Bauersachs R, Bellmunt-Montoya S, Black SA, ve ark. Editor's Choice – European Society for Vascular Surgery (ESVS) 2021 Clinical Practice Guidelines on the Management of Venous Thrombosis. *Eur J Vasc Endovasc Surg.* 2021;61(1):9-82.
37. Bozkurt AK, editör. Periferik arter ve ven hastalıkları ulusal tedavi kılavuzu. İstanbul: Bayçınar Tıbbi Yayıncılık; 2021.
38. Ortel TL, Neumann I, Ageno W, Beyth R, Clark NP, Cuker A, ve ark. American Society of Hematology 2020 guidelines for management of venous thromboembolism: treatment of deep vein thrombosis and pulmonary embolism. *Blood Adv.* 2020;4:4693-738.
39. National Institute for Health and Care Excellence. Venous thromboembolic diseases: diagnosis, management and thrombophilia testing (clinical guideline 144). Available at: www.nice.org.uk/guidance/ng158.
40. Tran HA, Gibbs H, Merriman E, Curnow JL, Young L, Bennett A, ve ark. New guidelines from the Thrombosis and Haemostasis Society of Australia and New Zealand for the diagnosis and management of venous thromboembolism *Med J Aust.* 2019;210:227-35.