

KOZMETİK ENDİKASYONLARDA TOPİKAL RETİNOİDLER

6.
Bölüm

Aslı TATLIPARMAK¹
Zehra AŞİRAN SERDAR²

ÖZET

Retinoidler, nükleer reseptörlere bağlanarak etkilerini gösteren bir grup A vitamini türevinin genel ismidir. Günümüze kadar ortaya çıkan retinoidler 1., 2. ve 3. kuşak olarak alt gruplara kategorize edilmişlerdir. Ortaya çıktıkları ilk yıllarda akne tedavisinde geniş yer bulmuşlar, yıllar içinde foto yaşlanmadan akantozis nigrikansa kadar pek çok endikasyonda kullanımları gündeme gelmiştir. İrirtasyon, eritem, yanma ve deskuamasyon zaman zaman kullanımını kısıtlayan en önemli yan etkilerdir. Bu bölümde, topikal retinoidlerin kozmetik endikasyonlarda kullanımından bahsedilecektir.

Giriş

Vitamin A (all trans retinol) ve onun doğal ve sentetik türevleri retinoidler olarak isimlendirilmektedir (1-3). Dermatolojide retinoidler, ilk olarak akne vulgarisde kullanılmış daha sonra bu endikasyona foto-yaşlanma, solar lentigo ve hiperpigmentasyon gibi endikasyonlar da eklenmiştir (2).

1. Foto Yaşlanması

Ekstrinsik yaşlanmadaki en önemli etkenlerden olan ultraviyole radyasyon (UVR)'a uzun süreli maruziyet pigmentasyon bozukluğu, elastikiyet kaybı, telenjiyekta ziler, dispigmentasyon ve kırışıklıklara yol açmaktadır (2,4).

Topikal retinoidlerin fotoyaşlanması ilk kullanımları, akne tanısı ile tretino-in krem kullanan hastaların periorbital bölge kırışıklıklarında azalma tariflemesi

¹ Doç Dr. Aslı TATLIPARMAK, İstinye Üniversitesi Tip Fakültesi Deri ve Zührevi Hastalıklar Bölümü aslitatliparmak@gmail.com

² Prof. Dr. Zehra AŞİRAN SERDAR İstanbul Yeniyüzyıl Üniversitesi Deri ve Zührevi Hastalıklar Bölümü zehra.serdar@yeniyuziyil.edu.tr

Kaynaklar

- Beckenbach L, Baron JM, Merk HF, Löfller H, Amann PM. Retinoid treatment of skin diseases [Internet]. Vol. 25, European Journal of Dermatology. John Libbey Eurotext; 2015. p. 384–91.
- Riahi RR, Bush AE, Cohen PR. Topical Retinoids: Therapeutic Mechanisms in the Treatment of Photodamaged Skin. *Am J Clin Dermatol.* 2016;17(3):265–76.
- Bulengo-Ransby SM, Griffiths CE, Kimbrough-Green CK, Finkel LJ, Hamilton TA, Ellis CN, et al. Topical tretinoin (retinoic acid) therapy for hyperpigmented lesions caused by inflammation of the skin in black patients. *N Engl J Med.* 1993; 328(20):1438–43.
- Lowe NJ, Gifford M, Tanghetti E, Poulin YP, Goldman M, Tse Y, et al. Tazarotene 0.1% cream versus tretinoin 0.05% emollient cream in the treatment of photodamaged facial skin: a multicenter, double-blind, randomized, parallel-group study. *J Cosmet Laser Ther.* 2004;6(2):79–85.
- Sunder S. Relevant Topical Skin Care Products for Prevention and Treatment of Aging Skin [Internet]. Vol. 27, Facial Plastic Surgery Clinics of North America. W.B. Saunders; 2019. p. 413–8.
- Zasada M, Budzisz E. Randomized parallel control trial checking the efficacy and impact of two concentrations of retinol in the original formula on the aging skin condition: Pilot study. *J Cosmet Dermatol.* 2020;19(2):437–43.
- Antoniou C, Kosmadaki MG, Stratigos AJ, Katsambas AD. Photoaging. *Am J Clin Dermatol.* 2010;11(2):95–102.
- Katsambas AD, Katoulis AC. Topical Retinoids in the Treatment of Aging of the Skin. In: Advances in Experimental Medicine and Biology. Kluwer Academic/Plenum Publishers; 1999. p. 477–82.
- Maddin S, Lauharanta J, Agache P, Burrows L, Zultak M, Bulger L. Isotretinoin improves the appearance of photodamaged skin: Results of a 36-week, multicenter, double-blind, placebo-controlled trial. *J Am Acad Dermatol.* 2000;42(1):56–63.
- Manaloto RMP, Alster TS. Periorbital Rejuvenation: A Review of Dermatologic Treatments. *Dermatologic Surg.* 1999;25(1):1–9.
- Kang HY, Valerio L, Bahadoran P, Ortonne J-P. The Role of Topical Retinoids in the Treatment of Pigmentary Disorders. *Am J Clin Dermatol.* 2009;10(4):251–60.
- Sheth VM, Pandya AG. Melasma: A comprehensive update. *J Am Acad Dermatol.* 2011;65(4):699–714.
- Griffiths CEM, Finkel LJ, Ditre CM, Hamilton TA, Ellis CN, Voorhees JJ. Topical tretinoin (retinoic acid) improves melasma. A vehicle-controlled, clinical trial. *Br J Dermatol.* 1993;129(4):415–21.
- Leenutaphong V, Nettakul A, Rattanasuwon P. Topical Izotretinoin for melasma in Thai patients: a vehicle-controlled clinical trial. *J Med Assoc Thai.* 1999;82(9):868–75.
- Dogra S, Kanwar AJ, Parsad D. Adapalene in the treatment of melasma: A preliminary report [1] [Internet]. Vol. 29, Journal of Dermatology. Japanese Dermatological Association; 2002. p. 539–40.
- Rafal ES, Griffiths CEM, Ditre CM, Finkel LJ, Hamilton TA, Ellis CN, et al. Topical Tretinoin (Retinoic Acid) Treatment for Liver Spots Associated with Photodamage. *N*

- Engl J Med. 1992;326(6):368–74.
- 17. Kang S, Goldfarb MT, Weiss JS, Metz RD, Hamilton TA, Voorhees JJ, et al. Assessment of adapalene gel for the treatment of actinic keratoses and lentigines: A randomized trial. *J Am Acad Dermatol.* 2003;49(1):83–90.
 - 18. Draeles ZD. The combination of 2% 4-hydroxyanisole (mequinol) and 0.01% tretinoin effectively improves the appearance of solar lentigines in ethnic groups. *J Cosmet Dermatol.* 2006;5(3):239–44.
 - 19. Ortonne JP, Camacho F, Wainwright N, Bergfelt L, Westerhof W, Roseeuw D. Safety and efficacy of combined use of 4-hydroxyanisole (mequinol) 2%/tretinoin 0.01% solution and sunscreen in solar lentigines. *Cutis.* 2004;74(4):261–4.
 - 20. Bulengo-Ransby SM, Griffiths CE, Kimbrough-Green CK, Finkel LJ, Hamilton TA, Ellis CN, et al. Topical tretinoin (retinoic acid) therapy for hyperpigmented lesions caused by inflammation of the skin in black patients. *N Engl J Med.* 1993;328(20):1438–43.
 - 21. Grimes P, Callender V. Tazarotene cream for postinflammatory hyperpigmentation and acne vulgaris in darker skin: a double-blind, randomized, vehicle-controlled study. *Cutis.* 2006;77(1):45–50.
 - 22. Zurada JM, Kriegel D, Davis IC. Topical treatments for hypertrophic scars. *J Am Acad Dermatol.* 2006;55(6):1024–31.
 - 23. Massaki ABMN, Fabi SG, Fitzpatrick R. Repigmentation of Hypopigmented Scars Using an Erbium-Doped 1,550-nm Fractionated Laser and Topical Bimatoprost. *Dermatologic Surg.* 2012;38(7pt1):995–1001.
 - 24. Treesirichod A, Chaithirayanon S, Wongjirat N. Comparison of the efficacy and safety of 0.1% adapalene gel and 0.025% tretinoin cream in the treatment of childhood acanthosis nigricans. *Pediatr Dermatol.* 2019;36(3):330–4.
 - 25. Tian T, Luo Y-J, Wang H, Chen H-D, Li Y-H. Efficacy and Safety of a Sublative Bipolar Fractional Radiofrequency System Combined With Topical Tretinoin in Treating Striae Gravidarum. *Dermatologic Surg.* 2019;45(10):1245–52.
 - 26. Sorg O, Antille C, Kaya G, Saurat JH. Retinoids in cosmeceuticals [Internet]. Vol. 19, *Dermatologic Therapy.* Dermatol Ther; 2006. p. 289–96.
 - 27. Saurat JH, Sorg O. Topical natural retinoids. The “proligand-non-ligand” concept. *Dermatology.* 1999;199 Suppl 1:1–2.
 - 28. Babamiri K, Nassab R. Cosmeceuticals: The evidence behind the retinoids. *Aesthetic Surg J.* 2010;30(1):74–7.
 - 29. Green C, Orchard G, Cerio R, Hawk JLM. A clinicopathological study of the effects of topical retinyl propionate cream in skin photoageing. *Clin Exp Dermatol.* 1998;23(4):162–7.
 - 30. Creidi P, Vienne MP, Ochonisky S, Lauze C, Turlier V, Lagarde M, et al. Profilometric evaluation of photodamage after topical retinaldehyde and retinoic acid treatment. *J Am Acad Dermatol.* 1998;39(6):960–5.
 - 31. Gupta AK, Gover MD, Nouri K, Taylor S. The treatment of melasma: A review of clinical trials. *J Am Acad Dermatol.* 2006;55(6):1048–65.
 - 32. Torras H. Retinoids in aging. *Clin Dermatol.* 1996;14(2):207–15.