

ÜÇ BOYUTLU
GÖRÜNTÜLEMENİN
ORTODONTİDE
KULLANIMI

Dr Öğr Üyesi Özüm DAŞDEMİR ÖZKAN

© Copyright 2020

Bu kitabın, basım, yayın ve satış hakları Akademisyen Kitabevi A.Ş.'ne aittir. Anılan kuruluşun izni alınmadan kitabın tümü ya da bölümleri mekanik, elektronik, fotokopi, manyetik kağıt ve/veya başka yöntemlerle çoğaltılamaz, basılamaz, dağıtılamaz. Tablo, şekil ve grafikler izin alınmadan, ticari amaçlı kullanılamaz. Bu kitap T.C. Kültür Bakanlığı bandrolü ile satılmaktadır.

ISBN

978-625-7106-84-9

Kitap Adı

Üç Boyutlu Görüntülemenin Ortodontide Kullanımı

Yazar

Dr Öğr Üyesi Özüm DAŞDEMİR ÖZKAN

Yayın Koordinatörü

Yasin Dilmen

Sayfa ve Kapak Tasarımı

Akademisyen Dizgi Ünitesi

Yayıncı Sertifika No

47518

Baskı ve Cilt

Bizim Dijital Matbaa

Bisac Code

MED016030

DOI

10.37609/akya.86

GENEL DAĞITIM

Akademisyen Kitabevi A.Ş.

Halk Sokak 5 / A

Yenişehir / Ankara

Tel: 0312 431 16 33

siparis@akademisyen.com

www.akademisyen.com

ÖNSÖZ

Gelişen teknoloji ile birlikte hayatımıza giren üç boyutlu görüntüleme birkaç on yıl içinde sadece ileri değerlendirilmede başvurulan bir teknik olmaktan ziyade temel görüntüleme yöntemi olarak diş hekimliğinde ki yerini alacaktır. Bu teknolojik gelişime ayak uydurmak tüm uzman diş hekimleri, diş hekimleri, lisans ve uzmanlık öğrencileri için gereklidir.

Bu kitapta üç boyutlu görüntüleme teknikleri, üçüncü boyutun ortodonti pratiğinde başvurulduğu durumlar genel başlıklar halinde ele alınmıştır. Üç boyutlu veriler üzerindeki sefalometrik değerlendirmeler, anatomik noktalar, hava yolu analizi detaylı olarak anlatılmış, görseller ile açıklanmıştır.

Özellikle üçüncü boyutu rutin olarak kullanacak, teknolojiyi yakından takip eden genç klinisyenlere pratik bir rehber kitap olarak faydalı olması dileklerle...

*“Verdiği Destek İçin Modern Ortodonti Firmasına
Teşekkür Ederiz”*

İÇİNDEKİLER

BÖLÜM 1

ORTODONTİDE GÖRÜNTÜLEME	1
Ortodontide Üç Boyutlu Görüntülemenin Tarihsel Gelişimi	3
İki Boyutlu Görüntüleme Teknikleri	4
Sefalometri ve Tarihiçesi.....	4
Geleneksel Sefalometrik Radyografi	5
Dijital Sefalometrik Radyografi	6
Ortodontide Geleneksel İki Boyutlu Sefalometrik Filmlerin Kullanım Alanları ve Kısıtlamaları	7
Kullanım Alanları	7
Kısıtlamaları	8
Lateral Sefalometrik Analiz Yöntemleri.....	11
Geleneksel Sefalometrik Analiz Yöntemi	11
Bilgisayarlı Sefalometrik Analiz Yöntemleri	11
Bilgisayarlı Sefalometrik Analiz Yazılımları.....	12
Ortodontide Üç Boyutlu Görüntülemeye Neden İhtiyaç Vardır?.....	13
Üç Boyutlu Görüntüleme Teknikleri	14
Manyetik Rezonans Görüntüleme (MRG)	14
Bilgisayarlı Tomografi (BT)	15
Konik Işınlı Bilgisayarlı Tomografi (Cone Beam Computed Tomography = CBCT):	17

Üç Boyutlu Görüntülemenin Ortodontide Kullanım Alanları...	21
Gömülü Kaninlerin Görüntülenmesi	21
Kök Rezorpsiyonunun Değerlendirilmesi	21
Kök Fraktürlerinin Değerlendirilmesi	21
Ortodontik Mini Vida Yerleştirilmesi	22
Asimetrinin Değerlendirilmesi	22
Temporomandibular Eklemdeki Dejeneratif Değişimlerin Değerlendirilmesi	22
Dudak-Damak Yarıklarının Değerlendirilmesi	23
Solunum Yolunun Değerlendirilmesi	23
Ortodontik Sanal Modelleme	23
Kraniyofasiyal Anomalilerin Değerlendirilmesi.....	23
Büyüme Gelişimin Değerlendirilmesi	24
Üç Boyutlu Sefalometrik Değerlendirme	24

BÖLÜM 2

ÜÇ BOYUTLU ANALİZ PROGRAMI İLE ANALİZ AŞAMALARI....29

Üç Boyutlu Verilerinin Elde Edilmesi.....	29
Üç Boyutlu Verilerinin Sefalometrik Analiz Programlarına Aktarılması ve Oryantasyonu	29
Üç Boyutlu Görüntüler Üzerindeki Anatomik İşaret Noktaları	35
Üç Boyutlu Hava Yolu Analizi	44

Kaynaklar49

Özgeçmiş67

İndeks69

KAYNAKLAR

- ADAMS GL, GANSKY SA, MILLER AJ, HARRELL WE, Jr H, DC. (2004) Comparison between traditional cephalometry and a 3- dimensional approach on human dry skulls. *Am J Orthod Dentofacial Orthop*,4.
- AHLQVIST J, ELIÄSSON S, WELANDER U. (1986) The effect of projection errors on cephalometric length measurements. *Eur J Orthod*, 8, 141-148.
- AHLQVIST J, ELIÄSSON S, WELANDER U. (1988) The effect of projection errors on angular measurements in cephalometry. *Eur J Orthod*, 10, 353-361.
- AJAYI E. O. (2005) Cephalometric norms of Nigerian children. *Am. J. Orthod. Dentofacial Orthop*, 128, 653-656.
- AKIN M, TEZCAN M, İLERİ Z. (2014) Konvansiyonel ve dijital sefalometrik ölçüm yöntemlerinin karşılaştırılması Comparison of conventional and digital cephalometric measurement methods. *SDÜ Sağlık Bilimleri Enstitüsü Dergisi Cilt 5 / Sayı 3*
- Van AKEN J, Van Der LINDEN LW. (1966) The integral absorbed dose in conventional and panoramic complete-mouth examinations. *Oral Surg Oral Med Oral Pathol*, 22, 603-616.
- AKGÜL AA, TOYGAR TU. (2002) Natural craniofacial changes in the third decade of life: a longitudinal study. *Am. J. Orthod. Dentofacial Orthop*, 122, 512-22.
- ALLEN W. (1963) Historical aspects of roentgenographic sefalometry. *Am J Orthod*,49,451-458. Alınmıştır: Uzel İ, Enacar A. Ortodontide Sefalometri (II. Baskı), Çukurova Üniversitesi Basımevi, Adana, 3-18.

- ALTINSEL H. (1961) Sagittal sefalometrik radiografilerde kullanılan teşhis metotlarının baz metodu ile mukayese ve kritiği uzmanlık tezi, İstanbul Üniversitesi Tıp Fakültesi Dış Hekimliği Okulu. Alınmıştır: UZEL İ, ENACAR A. (2000). Ortodontide Sefalometri. Çukurova Üniversitesi Basımevi, Adana.
- ARAI Y, TAMMISALO E, IWAI K, HASHIMOTO K, SHINODA K. (1999) Development of a compact computed tomographic apparatus for dental use. *Dentomaxillofac Radiol*, 28, 245-248
- BABA R, UEDA K, KUBA A, KOHDA E, SHIGARA N. (1999) Development of a subject-standing-type cone-beam CT for chest and orthopedic imaging. *Radiology*, 213, 194.
- BAILEY K, TAYLOR W. (1998) Mesh diagram cephalometric norms for Americans of African descent. *Am. J. Orthod. Dentofacial Orthop*, 114, 21822.
- BATURAY T, ERDOĞAN E. (1977) Normal okluzyonlu toplumumuz bireylerinde tweed normları. Doktora tezi. T. C. Gülhane Askeri Tıp Akademisi, Ankara.
- BAUMRIND S, FRANTZ R. (1971) The reliability of head film measurements. *Am J Orthod*, 60.
- BAUMRIND S. (2001a) Integrated three-dimensional craniofacial mapping: Background, principles, and perspectives. *Semin Ortho*, 4.
- BAUMRIND S. (2001b) Integrated three-dimensional craniofacial mapping: Background, principles, and perspectives. *Semin Orthod*, 4, 223-232.
- BAYDAŞ B, ERDEM A, YAVUZ İ, CEYLAN İ. (2007) Heritability of facial proportions and soft-tissue profile characteristics in Turkish Anatolian siblings. *Am. J. Orthod. Dentofacial Orthop*, 131, 504-513.
- BAYOME M, PARK JH, KOOK YA. (2013) New three-dimensional cephalometric analyses among adults with a skeletal Class I pattern and normal occlusion. *Korean J Orthod*, 43, 62-73.
- BAZ O. (1956) Sagittal anomalilerin uzak röntgen resimlerinin tetkik ve teşhisi. Doçentlik Tezi. Alınmıştır: UZEL İ, ENACAR A. (2000). Ortodontide Sefalometri. Adana: Çukurova Üniversitesi Basımevi.
- BEHBEHANI F, HICKS EP, BEEMAN C, KLUEMPER GT, RAYENS MK. (2006) Racial variations in cephalometric analysis between Whites and Kuwaitis. *Angle Orthod*, 76, 406-11.
- BEN-BASSAT Y, DINTE A, BRIN I, KOYOUNDJISKY-KAYE E. (1992) Cephalometric pattern of Jewish East European adolescents with clinically acceptable occlusion. *Am. J. Orthod. Dentofacial Orthop*, 102, 443-8.

- BERCO M, RIGALI PH Jr, MİNER RM, DELUCA S, ANDERSON NK, WILL LA. (2009) Accuracy and reliability of linear cephalometric measurements from cone-beam computed tomography scans of a dry human skull. *Am J Orthod Dentofacial Orthop*, 136, 17-19.
- BHOLSITHI W, THARANON W, CHINTAKANON K, KOMOLPIS R, SINTHANAYOTHIN C. (2009) 3D vs. 2D cephalometric analysis comparisons with repeated measurements from 20 Thai males and 20 Thai females. *Biomed Imaging Interv J*, 5,21.
- BIANCHI S, LOJACONO A. (1998) 2D and 3D images generated by Cone Beam Computed Tomography (CBCT) for dentomaxillofacial investigations. *Computer Assisted Radiology and Surgery*, 79, 2-7.
- BOLLEN AM. (2004) Three-dimensional scans and radiation exposure. *Am J Orthod Dentofacial Orthop*, 126, 17.
- BORG E, GRONDAHL HG. (1996) On the dynamic range of different X-ray photon detectors in intra-oral radiography. A comparison of image quality in film, charge-coupled device and storage phosphor systems. *Dentomaxillofac Radiol*, 28, 82-88.
- BRANNAN J. (2002) An introduction to dijital radiography in dentistry. *J Orthod*. 29:66-69
- BROADBENT BH. (1931) A new x-ray technique and its aplication to orthodontia. *Angle Orthod*, 2, 45-66.
- BRUNTZ LQ, PALOMO JM, BADEN S, HANS MG. (2006) A comparison of scanned lateral cephalograms with corresponding original radiographs. *Am J Orthod Dentofacial Orthop*, 130, 340-348.
- BURKE PH, BEARD FH. (1967) Stereophotogrammetry of the face. A preliminary investigation into the accuracy of a simplified system evolved for contour mapping by photography. *Am J Orthod*, 53, 769-782.
- BÜYÜK SK, RAMOĞLU Sİ. (2011) Cone Beam Computed Tomography in Orthodontic Diagnosis. *Sağlık Bilimleri Dergisi (Journal of Health Sciences)* 20(3) 227-234).
- CATTANEO PM, BLOCH CB, CALMAR D, HJORTSHOJ M, MELSEN B. (2008) Comparison between conventional and cone-beam computed tomography-generated cephalograms. *Am J Orthod Dentofacial Orthop*, 134, 798-802.
- CEVIDANES LH, FRANCO AA, GERIG G et al. (2005a) Comparison of relative mandibular growth vectors with high-resolution 3-dimensional imaging. *Am J Orthod Dentofacial Orthop*, 128, 27-34.
- CEVIDANES LH, FRANCO AA, GERIG G et al. (2005b) Assessment of mandibular growth and response to orthopedic treatment with

- 3-dimensional magnetic resonance images. *Am J Orthod Dentofacial Orthop*, 128, 16-26.
- CEYLAN İ, GAZİLERLİ Ü. (1992). Erzurum yöresi çocuklarındaki Steiner, Downs ve Tweed ölçümlerinin diğer bazı gruplarla karşılaştırılması. *Ankara Üniversitesi Diş Hekimliği Fakültesi Dergisi*, 19, 143-152.
- CHEESMAN K (2008) Anatomy of the naso-and oropharynx, Anaesthesia and Intensive Care Medicine, 9, 277-279.
- CHEN Y, HONG Y, WU K, CHEN M, CHAN H, CHEN K. (1993) Jaw triangle analysis: an adjuvan diagnostics. *Chin Dent J*, 12, 56-70.
- CHEN YJ, CHEN SK, HUANG HW, YAO CC, CHANG HF. (2004) Reliability of landmark identification in cephalometric radiography acquired by a storage phosphor imaging system. *Dentomaxillofac Radiol*, 33, 301-306.
- CHEN YJ, CHEN SK, YAO JC, CHANG HF. (2004) The effects of differences in landmark identification on the cephalometric measurements in traditional versus digitized cephalometry. *Angle Orthod*, 74, 155-161.
- CHIEN PC, PARKS ET, ERASO F, HARTSFİELD JK, ROBERTS WE, OFNER S. (2009) Comparison of reliability in anatomical landmark identification using two-dimensional digital cephalometrics and three-dimensional cone beam computed tomography in vivo. *Dentomaxillofac Radiol*, 38, 262-273.
- CHEUNG LK, CHAN YM, JAYARATNE YS, LO J. (2011) Threedimensional cephalometric norms of Chinese adults in Hong Kong with balanced facial profile. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 112:e56-73.
- CİĞER S. (1980) Ankara ve yöresinde normal kapanışlı genç erişkinlerin yumuşak doku profillerinin incelenmesi. *Hacettepe Üniversitesi Diş Hekimliği Fakültesi Dergisi*, 4, 69-77.
- COOKE S, WEI SHY. (1987) A comparative study of southern Chinese and British Caucasian cephalometric standards. *Angle Orthod*, 59, 131-138.
- COTTON WN, TAKANO WS, WONG WW. (1951) The downs analysis applied to three other ethnic groups. *Angle Orthod*, 21, 213-20.
- CUTTING C, GRAYSON B, BOOKSTEIN F, FELLINGHAM L, MCCARTHY JG. (1986) Computer-aided planning and evaluation of facial and orthognathic surgery. *Clin Plast Surg*, 13, 449-462.
- ÇIRAK F, ORHAN M, ÜŞÜMEZ S, GÜRAY E. (2002) Türk ortodontistlerinin tanı ve tedavi yaklaşımları. *Türk Ortodonti Dergisi*, 15, 8-14.

- DANFORTH R, DUS I, MAH J. (2003) 3-D volume imaging for dentistry: A new dimension. *CDA Journal*, 31, 17–23.
- DEVANNA, R. (2015) Two-dimensional to three-dimensional: A new three-dimensional cone-beam computed tomography cephalometric analysis. *Journal of Orthodontic Research* 3.1,30.
- DOLPHIN IMAGING AND MANAGEMENT SOLUTION. (2011). Dolphin Imaging User's Guide. Retrieved from: <http://www.promed.ua/wp-content/uploads/Dolphin%20full.pdf>
- DUNN GF, GREEN LJ, CUNAT JJ. (1973) Relationships between variation of mandibular morphology and variation of nasopharyngeal airway size in monozygotic twins. *Angle Orthod.* 43:129–135.
- DULA K, MINI R, Van Der STELT PF, LAMBRECHT JT, SCHNEEBERGER P, BUSER D. (1996) Hypothetical mortality risk associated with spiral computed tomography of the maxilla and mandible. *Eur J Oral Sci*, 104, 503-510.
- EKESTUBBE A, THILANDER-KLANG A, LITH A, GRONDAHL HG. (2004) Effective and organ doses from scanography and zonography: a comparison with periapical radiography. *Dentomaxillofac Radiol*, 33, 87-92.
- ELEFTERIADIS J, ATHANASIOU A. (1996) Evaluation of impacted canines by means of computerized tomography. *Int J Adult Orthodon Orthognath Surg*, 11(3), 257–64.
- ENLOW DH, HANS MG. (1996) Essentials of facial growth. Philadelphia: W.B. Saunders, p. 72.
- ERBAY EF, CANIKLIOĞLU CM, ERBAY SK. (2002) Soft tissue profile in Anatolian Turkish adults: Part I. Evaluation of horizontal lip position using different soft tissue analyses. *Am. J. Orthod. Dentofacial Orthop*, 121, 57-64.
- ERBAY EF, CANIKLIOĞLU CM. (2002) Soft tissue profile in Anatolian Turkish adults: Part II. Comparison of different soft tissue analyses in the evaluation of beauty. *Am. J. Orthod. Dentofacial Orthop.* 121: 65-72.
- ERKAN M, GÜREL HG, NUR M, DEMİREL B. (2011) Reliability of four different computerized cephalometric analysis programs. *Eur J Orthod* Apr 18, 2011 [Epub ahead of print].
- ERICSON S, KUROL J. (1988) Early treatment of palatally erupting maxillary canines by extraction of the primary canine. *Eur J Orthod*; 10(4): 283–95.

- EVANKO AM, FREEMAN KC, GEORGE J. (1996) Mesh diagram analysis: Developing a norm for Puerto Rican Americans. *Angle Orthod.* 67: 381-388.
- FRANCHI L, BACCETTI T, MCNAMARA JA Jr. (1998) Cephalometric floating norms for North American adults. *Angle Orthod.* 68: 497-502.
- FRAENKEL R. (1980) The applicability of the occipital reference base in cephalometrics. *Am J Orthod*; 77(4): 379-395.
- FREDERIKSEN NL, BENSON BW, SOKOLOWSKI TW. (1995) Effective dose and risk assessment from computed tomography of the maxillofacial complex. *Dentomaxillofac Radiol*; 24: 55-58.
- FUHRMANN RAW. (2002) Three-dimensional cephalometry and three-dimensional skull models in orthodontic/surgical diagnosis and treatment planning. *Semin Orthod*; 1.
- FUHRMANN RAW. (2002) Three-dimensional evaluation of periodontal remodeling during orthodontic treatment. *Semin Orthod*; 1.
- GAZİLERLİ Ü. (1976) Normal kapanışlı 13-16 yaşlar arasındaki Ankara çocuklarında Steiner normları. Doçentlik tezi. T. C. Ankara Üniversitesi Diş Hekimliği Fakültesi, Ankara.
- GAZİLERLİ Ü. (1981) Türk Çocukları için Downs ve Tweed ölçümleri. *Ankara Üniversitesi Diş Hekimliği Fakültesi Dergisi*, 8:115-136
- GAZİLERLİ Ü. (1982) Değişik cins ve yaş gruplarında Ricketts yumuşak doku ölçümleri. *Ankara Üniversitesi Diş Hekimliği Fakültesi Dergisi*. 9: 15-22.
- GAZİLERLİ Ü, CEYLAN İ. (1995) ANB açısını etkileyen etkenlerin incelenmesi. *Türk Ortodonti Dergisi*, 8, 182-186.
- GEIST JR, BRAND JW. (2001) Sensitometric comparison of speed group E and F dental radiographic films. *Dentomaxillofac Radiol*, 30, 147-152.
- GELGÖR İE, KARAMAN AI, ZEKİÇ E. (2006) The use of parental data to evaluate soft tissues in an Anatolian Turkish population according to Holdaway soft tissue norms. *Am. J. Orthod. Dentofacial Orthop*, 3, 330-9.
- GRAVELY JF, BENZİES PM. (1984) The clinical significance of tracing error in cephalometry. *Br J Orthod*, 11, 44-48.
- GRIBEL BF, GRIBEL MN, FRAZAO DC, MCNAMARA JA, MANZI FR. (2011) Accuracy and reliability of craniometric measurements on lateral cephalometry and 3D measurements on CBCT scans. *Angle Orthod*, 81, 26-35.

- GRUBB JE, SMITH T, SINCLAIR P.(1996) Clinical and scientific applications/advances in video imaging. *Angle Orthod*, 66, 407-416.
- GOLDSMAN S. (1959) The variations in skeletal and denture patterns in excellent adult facial types. *Angle Orthod*, 29, 63-92.
- GOTTLIEB EL, NELSON AH, VOGELS DS. (1996) JCO Study of Orthodontic Diagnosis and treatment procedures. Part 2. Breakdowns of selected variables. *J Clin Orthod*, 30, 689-698.
- GÜLYURT M. (1989) Erzurum yöresi çocuklarında Ricketts' in frontal sefalometrik ölçümleri. *Türk Ortodonti Dergisi*, 2, 144-151.
- GÜRSOY N, UĞUR T, GÜRSOY S. (1973). Normal kapanışlı Türklerde sefalometrik norm araştırması. *İstanbul Üniversitesi Diş Hekimliği Fakültesi Dergisi*,7, 333-343.
- HAGEMANN K, VOLLMER D, NIEGEL T, EHMER U, REUTER I. (2000) Prospective study on the reproducibility of cephalometric landmarks on conventional and digital lateral headfilms. *J Orofac Orthop*, 61, 91-99.
- HAJEER MY, MILLETT DT, AYOUB AF, SIEBERT JP. (2004) Applications of 3D imaging in orthodontics: part I. *J Orthod*, 31, 62-70.
- HAJIGHADIMI M, DOUGHERTY HL, GRAKANI F. (1981) Cephalometric evaluation of Iranian children and its comparasion with Tweed's and Steiner's standards. *Am. J. Orthod*. 79, 192-197.
- HALAZONETIS DJ. (2005) From 2-dimensional cephalograms to 3-dimensional computed tomography scans. *Am J Orthod Dentofacial Orthop*, 127, 627-637.
- HARDIN WJ. (1971) A cephalometric supplement for clinical application. *Am. J. Orthod*, 59, 568-580.
- HARRELL WE, Jr, HATCHER DC, BOLT RL. (2002) In search of anatomic truth: 3-dimensional digital modeling and the future of orthodontics. *Am J Orthod Dentofacial Orthop*, 122, 325-330.
- HARRELL WE. Jr. (2004) Planning: The future of three-dimensional analysis-Part1. *SAO News*, Fall.
- HASSAN B, Van Der STELT P, SANDERINK G. (2009) Accuracy of three-dimensional measurements obtained from cone beam computed tomography surface-rendered images for cephalometric analysis: influence of patient scanning position. *Eur J Orthod*, 31, 129-134.
- HASSAN AH. (2006) Cephalometric norms for Saudi adults living in the western region of Saudi Arabia. *Angle Orthod*, 76, 109-113.
- HATCHER DC, ABOUDARA CL. (2004) Diagnosis goes digital. *Am J Orthod Dentofacial Orthop*, 125, 512-515.

- HERRING J. (2007) The effectiveness of orthodontists and oral radiologists in the diagnosis of impacted maxillary canines. *Am J Orthod Dentofacial Orthop*, 132(6), 861.
- HIDEKI K, KOICHI Y, KAZUhide K, TAKASASHI N. (2000) Standardization of 3-D CT measurements for length and angles by matrix transformation in the 3-D coordinate system. *Cleft Palate Craniofacial J*, 37, 349-56.
- HONEY O, SCARFE W, HILGERS M. (2007) Accuracy of the cone-beam computed tomography imaging of the temporomandibular joint: comparisons with panoramic radiology and linear tomography. *Am J Orthod Dentofacial Orthop*, 132(4), 429-38.
- HOUSTON W. (1983) The analysis of errors in orthodontic measurements. *Am J Orthod*, 83, 382 - 390.
- HOUSTON W, MAHER R, MCELROY D, SHERRIFF M. (1986) Sources of error in measurements from cephalometric radiographs. *Eur J Orthod*, 8, 149-151.
- HUNTJENS E, KISS G, WOUTERS C, CARELS C. (2008) Condylar asymmetry in children with juvenile idiopathic arthritis assessed by cone-beam computed tomography. *Eur J Orthod*, 30, 545-51.
- HUANG WJ, TAYLOR RW, DASANAYAKE AP. (1998) Determining cephalometric norms for Caucasians and African Americans in Birmingham. *Angle Orthod*, 68, 503-11.
- HWANG HS, KIM WS, MCNAMARA JA. Jr. (2002) Ethnic differences in the soft tissue profile of Korean and European-American adults with normal occlusions and well-balanced faces. *Angle Orthod*, 72, 72-80.
- HWANG HS, YOUN IS, LEE KH, LIM HJ. (2007) Classification of facial asymmetry by cluster analysis. *Am J Orthod Dentofacial Orthop*, 132, 271-276.
- İŞIKSAL E. (1989) Normal kapanış ve dengeli yüz yapısı gösteren bireylerde steiner ölçüm değerleri. *Ege Üniversitesi Diş Hekimliği Fakültesi Dergisi*, 10, 718.
- İŞİMER Y, UZEL İ, SAĞDIÇ D. (1990) Toplumumuzdaki nötr okluzyonlu birey normlarının 124meri normları ile karşılaştırılması. *Türk Ortodonti Dergisi*, 3, 65-71.
- İŞCAN H. (1991) Ortodontik muayene ve ortodontik tedaviler, *Ankara: G.Ü. Diş Hek. Vakfı G.Ü.İ.V.yayımları*, 9-167.
- JACOBSON A, JACOBSON R. (2006) Radiographic cephalometry from basics to 3- D Imaging, 2nd ed.

- JAFFRAY DA, SIEWERDSEN JH. (2000) Cone-beam computed tomography with a flat-panel imager: initial performance characterization. *Med Phys*, 27, 1311-1323.
- JAKOBSONE G, NEIMANE L, KRUMINA G. (2010) Two and three dimensional evaluation of the upper airway after bimaxillary correction of class III malocclusion. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 110, 234-242.
- JANSON G, CRUZ KS, WOODSIDE DG, METAXAS A, de FREITAS MR, HENRIQUES JF. (2004) Dentoskeletal treatment changes in Class II subdivision malocclusions in submentovertex and postero-anterior radiographs. *Am J Orthod Dentofacial Orthop*, 126, 451-463.
- JOSSI V, YAMAGUCHI T, MATSUDA Y, KANEKO N, MAKI K, OKANO T. (2012) Skeletal maturity assessment with the use of cone-beam computerized tomography. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 113,(6):841-9
- KANTOR ML, PHILLIPS CL, PROFFIT WR. (1993) Subtraction radiography to assess reproducibility of patient positioning in cephalometrics. *Am J Orthod Dentofacial Orthop*, 104, 350-354.
- KAU CH, RICHMOND S, PALOMO JM, HANS MG. (2005) Three-dimensional cone beam computerized tomography in orthodontics. *J Orthod*, 32, 282-293.
- KAZANDJIAN S, KILIARIDIS S, MAVROPOULOS A. (2006) Validity and reliability of a new edge-based computerized method for identification of cephalometric landmarks. *Angle Orthod*, 76, 619-624.
- KIM EJ, PALOMO JM, KIM SS, LIM HJ, LEE KM, HWANG HS. (2011) Maxillofacial characteristics affecting chin deviation between mandibular retrusion and prognathism patients. *Angle Orthod*, 81, 988-93.
- KILIÇ N, ÇATAL G, OKTAY H. (2010) McNamara norms for Turkish adolescents with balanced faces and normal occlusion. *Aust Orthod J*. May, 26(1), 33-7.
- KITAURA H, YONETSU K, KITAMORI H, KOBAYASHI K, NAKAMURA T.(2000) Standardization of 3-D CT measurements for length and angles by matrix transformation in the 3-D coordinate system. *Cleft Palate Craniofac J*, 4, 349-356.
- KOCH S, WAGNER IV, SCHNEIDER W. (2000) Effective and quality-controlled use of digital radiography in dental practice. *Int J Comput Dent*, 3, 107-118.

- KOCADERELİ İ, TELLİ AE. (1999) Evaluation of Ricketts' long-range growth prediction in Turkish children. *Am. J. Orthod. Dentofacial Orthop*, 115, 515-20.
- KOWALSKI CJ, NASJLETI CE, WALKER GF. (1975) Dentofacial variations within and between four groups of adult 124merican males. *Angle Orthod*, 45, 146-151.
- KUMAR V, LUDLOW JB, MOL A, (2007) Cevidanes L. Comparison of conventional and cone beam CT synthesized cephalograms. *Dentomaxillofac Radiol*, 36, 263-269.
- KURODA S, YAMADA K, DEGUCHI T. (2007) Root proximity is a major factor for screw failure in orthodontic anchorage. *Am J Orthod Dentofacial Orthop*, 131, 68-73.
- LAGRAVERE MO, LOW C, FLORES-MIR C et al. (2005) Intraexaminer and interexaminer reliabilities of landmark identification on digitized lateral cephalograms and formatted 3-dimensional cone-beam computerized tomography images. *Am J Orthod Dentofacial Orthod*, 137, 598-604.
- LASCALA CA, PANELLA J, MARQUES MM. (2004) Analysis of the accuracy of linear measurements obtained by cone beam computed tomography (CBCT-NewTom). *Dentomaxillofac Radiol*, 33, 291-294.
- LEE CT. (1988) Standards for Korean adult facial relationships by various roentgeno-cephalometric analysis. *Korean J Orthod*, 18, 459-74.
- LEE H, BAYOME M, KIM SH, KIM KB, BEHRENTS RG, KOOK YA. (2012) Mandibular dimensions of subjects with asymmetric skeletal Class III malocclusion and normal occlusion compared with cone-beam computed tomography. *Am J Orthod Dentofacial Orthop*, 142, 179-85.
- LORENZONI DC, BOLOGNESE AM, GARIB DG, GUEDES FR, & SANT'ANNA EF. (2012). Cone-beam computed tomography and radiographs in dentistry: aspects related to radiation dose. *International journal of dentistry* 2012.
- LOUBELE M, BOGAERTS R, Van DIJCK E, PAUWELS R, VANHEUSDEN S, SUETENS P et al. (2009) Comparison between effective radiation dose of CBCT and MSCT scanners for dentomaxillofacial applications. *Eur J Radiol*, 71, 461-468.
- LUDLOW JB, IVANOVIC M. (2008) Comparative dosimetry of dental CBCT devices and 64-slice CT for oral and maxillofacial radiology. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 106, 106-114.

- LUDLOW JB, GUBLER M, CEVIDANES L, MOL A. (2009) Precision of cephalometric landmark identification: cone-beam computed tomography vs conventional cephalometric views. *Am J Orthod Dentofacial Orthop*, 136, 311-310.
- MACCHI A, CARRAFIELLO G, CACCIAFESTA V, NORCINI A. (2006) Three-dimensional digital modeling and setup. *Am J Orthod Dentofacial Orthop*, 129, 605-610.
- MAH J. (2007) The evolution of digital study models. *J Clin Orthod*, 41, 557-561.
- MAKI K, USUI T, KUBOTA M, NAKANO H, SHIBASAKI Y. (2002) Application of cone-beam X-ray CT in dento-maxillofacial region. *Computer Assisted Radiology and Surgery*, 3-8.
- MANSSON S, JOHANSSON E, MAGNUSSON P, CHAI CM, HANSSON G, PETERSSON JS et al. (2006) 13C imaging-a new diagnostic platform. *Eur Radiol*, 16, 57-67.
- MCINTYRE GT, MOSSEY PA. (2003) Size and shape measurement in contemporary cephalometrics. *Eur J Orthod*, 25, 231-242.
- MCNAMARA JA. (1984). A method of cephalometric evaluation. *Am J Orthod*, 86, 449-469.
- MIDTGARD J, BJORK G, LINDER-ARONSON S. (1974) Reproducibility of cephalometric landmarks and errors of measurements of cephalometric cranial distances. *Angle Orthod*, 44, 56-61.
- MIURA F, INOUE N, SUZUKI K. (1963) The standards of steiner's analysis for Japanese. *Bull. Tokyo Med. And Dent. Univ*, 10: 387-395.
- Alınmıştır: CEYLAN İ, GAZİLERLİ, Ü. (1992). *Ankara Üniversitesi Dış Hekimliği Fakültesi Dergisi*, 19, 143-152.
- MIRACLE AC, MUKHERJI SK. (2009) Conebeam CT of the head and neck, part 2: clinical applications. *AJNR Am J Neuroradiol*, 30, 1285-1292.
- de MORAES ME, HOLLENDER LG, CHEN CS, MORAES LC, BALDUCCI I. (2011) Evaluating craniofacial asymmetry with digital cephalometric images and cone-beam computed tomography. *Am J Orthod Dentofacial Orthop*, 139, 523-531.
- MIYAJIMA K, MCNAMARA JA Jr., KIMURA T, MURATA S, LIZUKA T. (1996) Craniofacial structure of Japanese and European-American adults with normal occlusions and well-balanced faces. *Am. J. Orthod. Dentofacial Orthop*, 110, 431-8.
- MOLDEZ MA, SATO K, SUGAWARA J, MITANI H. (2006). Linear and Angular Filipino Cephalometric Norms According to Age and Sex. *Angle Orthod*, 76, 800-805.

- MOREIRA CR, SALES MA, LOPES PM, CAVALCANTI MG. (2009) Assessment of linear and angular measurements on three-dimensional cone-beam computed tomographic images. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 108, 430-436.
- MORI Y, MIYAJIMA T, MINAMI K, SAKUDA M. (2001) An accurate three-dimensional cephalometric system: a solution for the correction of cephalic malpositioning. *J Orthod*, 28, 143-149.
- MOYER R, BOOKSTEIN F. (1979) The inappropriateness of conventional cephalometrics. *Am J Orthod*, 75, 599- 617.
- NAKAGAWA Y, KOBAYASHI K, ISHII H, MISHIMA A, ASADA K, ISHIBASHI K. (2002) Preoperative application of limited cone beam computerized tomography as an assessment tool before minor oral surgery. *Int J Oral Maxillofac Surg*, 31, 322-326.
- NAKASİMA A, TERJİMA M, MORİ N et al. (2005) Three-dimensional computer-generated head model reconstructed from cephalograms, facial photographs, and dental cast models. *Am J Orthod Dentofacial Orthop* ; 127: 282-292.
- NALÇACI R, ÖZTÜRK F, SÖKÜCÜ O. (2010) A comparison of two-dimensional radiography and three-dimensional computed tomography in angular cephalometric measurements. *Dentomaxillofac Radiol*, 39, 100-106.
- NEWTON TH, POTTS DG. (1981) Technical Aspects of Computed Tomography. *Radiology of the skull and brain*, 5, 3853-4419.
- NING R, KRUGER R. (1988) Computed simulation of image intensifier-based computed tomography detector: vascular application. *Phys Med Biol*, 15, 188-92.
- de OLIVEIRA AE, CEVIDANES LH, PHILLIPS C, MOTTA A, BURKE B, TYNDALL D. (2009) Observer reliability of three-dimensional cephalometric landmark identification on cone-beam computerized tomography. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 107, 256-265.
- ONKOSUWITO EM, KATSAROS C, VAN'T HOF M, BODEGOM J, KUIPERS-, JAGTMAN A. (2002) The reproducibility of cephalometric measurements: a comparison of analogue and digital methods. *Eur J Orthod*, 24, 655 – 665.
- OKTAY H. (1991) A comparison of ANB, WITS, AF-BF, APDI measurements. *Am. J. Orthod. Dentofacial Orthop*, 99, 122-8.
- ORHAN K. (2012) Diş Hekimliğinde konik ışıklı komputerize tomografinin yeri ve önemi. *Yeditepe Üniveristesi Diş Hekimliği Fakültesi Dergisi*, 3(3):6-17

- ORHAN K, AKSOY S. (2015) Konik Işınli Bilgisayalı Tomografi ile Üç Boyutlu Sefalometri (Bölüm 15) Güncel Bilgiler Işığında Ortodonti, E ÖZDİLER. 1. Baskı, Gümüş Kitabevi, Ankara, 295-360.
- ÖZ U, Orhan K, Abe N. (2011) Comprasion of lineer anguler measurements using two-dimensionel conventional methods and three-dimensionel cone beam CT images reconstructed from a volumetric rendering program in vivo. *Dentomaksillofac Radiol*,40:492-500
- ÖZTÜRK Y. (1983) Normal kapanışlı erişkin türklerin Björk'ün sefalometrik analiz normlarına göre incelenmesi. *İstanbul Üniversitesi Diş Hekimliği Fakültesi Dergisi*, 17,158-166.
- PACINI A. (1922) Roentgen ray anthropology of the skull. *J Radiol*, 3, 230-231, 322-331, 418-426.
- PAPADOPOULOS M, JANNOWITZ C, BOETTCHER P et al. (2000) Three-dimensional fetal cephalometry: An evaluation of the reliability of cephalometric measurements based on threedimensional CT reconstructions and on dry skulls of sheep fetuses. *Journal of Cranio-Maxillofacial Surgery*, 33, 229-237.
- PARK IC, BOWMAN D, KLAPPER L. (1989) A cephalometric study of Korean adults. *Am J Orthod Dentofacial Orthop*; 96: 5459.
- PECK J, SAMESHIMA G, MILLER A. (2007) Mesiodistal root angulation using panoramic and cone beam CT. *Angle Orthod*, 77(2), 206-13.
- POWER G, BRECKON J, SHERRIFF M, MCDONALD F. (2005) Dolphin Imaging Software: an analysis of the accuracy of cephalometric digitization and orthognathic prediction. *Int J Oral Maxillofac Surg*, 34, 619-626.
- PREDA L, La FIANZA A, Di MAGGIO EM et al. (1997) The use of spiral computed tomography in the localization of impacted maxillary canines. *Dentomaxillofac Radiol*, 26, 236-241.
- QUINTERO JC, TROSIEN A, HATCHER D, KAPILA S. (1999) Craniofacial imaging in orthodontics: historical perspective, current status, and future developments. *Angle Orthod*, 69, 491-506.
- RAKOSI T. (1982) Cephalometric radiography. London: Wolfe Medical Publications Ltd,7-223.
- RAPER H. (1953) Notes on early history of radiodontia, *Oral Surg* 1953; 6: 70- 81. Alınmıştır: Uzel İ, Enacar A. Ortodontide Sefalometri (II. Baskı), Çukurova Üniversitesi Basımevi, Adana,3-18.
- RICKETTS RM. (1960a). A foundation for cephalometric communication. *Am. J. Orthod*, 46, 330-57.

- RICKETTS RM. (1960b). The influence of orthodontic treatment on facial growth and development. *Angle Orthod*, 30, 103-133.
- RICKETTS RM. (1961) Cephalometric Analysis And Synthesis. *Angle Orthod*,3.
- RITMAN E, RITMAN E, ROBB R, GILBERT B, HARIS L. (1980) Three dimensional imaging of heart, lungs, and circulation. *Science*, 210, 273-80.
- ROBB RA. (1982) The Dynamic spatial reconstructor: An X-Ray Video-Fluoroscopic CT scanner for dynamic volume imaging of moving organs. *IEEE Trans Med Imaging*, 1, 22-33.
- SAINI S, SHARMA R, LEVINE LA, BARMSON RT, JORDAN PF, TH-RALL JH. (2001) Technical cost of CT examinations. *Radiology*, 218, 172-175.
- SAVARA BS, TRACY WE, MILLER PA. (1966) Analysis of errors in cephalometric measurements of three-dimensional distances on the mandible. *Arch Oral Biol*, 11, 209-217.
- SAYINSU K, İŞİK F, TRAKYALI G, ARUN T. (2007) An evaluation of the errors in cephalometric measurements on scanned cephalometric images and conventional tracings. *Eur J Orthod*, 29, 105-108.
- SCARFE WC, FARMAN AG, SUKOVIC P. (2006) Clinical applications of cone-beam computed tomography in dental practice. *J Can Dent Assoc*, 72, 75-80.
- SCARFE WC, FARMAN AG. (2008) What is cone-beam CT and how does it work? *Dent Clin North Am*,52,707-730.
- SCHWAB RJ (1998) Upper airway imaging, *Clin Chest Med*, 19, 33-54.
- SCHWARTZ R. (1927) Cephalometric method and Orthodontics. *J Am Dent Ass*, 14,22-35. Alınmıştır: Uzel İ, Enacar A. Ortodontide Se-falometri. II. Baskı, Çukurova Üniversitesi Basımevi, Adana, 2000, 3-18.
- SEKIHARA K, KAWAI H, YAMAMOTO K, KUMAZAKI T. (1995) Co-ne-beam CT angiography. *Proc JAMIT Front*, 33-8.
- SELVIK GA. (1974) Roentgen stereophotogrammetric method for the study of the kinematics of the skeletal systems. Malmö, Sweden: University of Lund.
- SİEWERDSEN JH, JAFFRAY DA. (1999) Cone-beam computed tomography with a flat-panel imager: *effects of image lag Med Phys*, 26, 2635-2647.
- SILVA MA, WOLF U, HEINICKE F, BUMANN A, VISSER H, HIRSCH E. (2008) Cone-beam computed tomography for routine orthodon-

- tic treatment planning: a radiation dose evaluation. *Am J Orthod Dentofacial Orthop*, 133, 640-645.
- SINGH IJ, SAVARA BS. (1966) Norms of size and annual increments of seven anatomical measures of maxillae in girls from three to sixteen years of age. *Angle Orthod*, 36, 312-324.
- SHAH SM, JOSHI MR. (1978) An assessment of asymmetry in the normal craniofacial complex. *Angle Orthod*, 48, 141-8.
- STABRUN AE, DANIELSEN K. (1982) Precision in cephalometric landmark identification. *Eur J Orthod*, 4, 185-196.
- STEINER CC. (1953) Cephalometrics for you and me. *Am. J. Orthod*, 39, 729-736.
- STEINER CC. (1959) Cephalometrics in clinical practice. *Angle Orthod*, 29, 8- 29.
- STEWART J, HEO G, GLOVER K. (2001) Factors that relate to treatment duration for patients with palatally impacted maxillary canines. *AmJ Orthod Dentofacial Orthop*, 119(3), 216-25.
- SWENNEN G, SCHUTYSER F, HAUSAMEN J. (2005) Three dimensional cephalometry. Springer Verlag, Berlin.
- SWENNEN GRJ, SCHUTYSER F, HAUSAMEN J-E. (2006) Chapter 3 - 3D cephalometric system. In: Three-dimensional cephalometry: a color atlas and manual. Berlin: Springer Verlag, 99-105.
- SWENNEN GR, SCHUTYSER F. (2006) Three-dimensional cephalometry: spiral multi-slice vs cone-beam computed tomography. *Am J Orthod Dentofacial Orthop*, 130, 410-416.
- SWLERENGA D, OESTERLE LJ, MESSERSMITH ML. (1994) Cephalometric values for adult Mexican-American. *American Journal of Orthodontics and Dentofacial Orthopedics*, 106(2), 146-155.
- THILANDER B, PERSSON M, ADOLFSSON U. (2005) Roentgen cephalometric standards for a Swedish population. A longitudinal study between the ages of 5 and 31 years. *Eur J Orthod*, 27, 370-89.
- TITİZ İ, LAUBINGER M, KELLER T, HERTRICH K, HIRSCHFELDER U. (2011) Repeatability and reproducibility of landmarks--a three-dimensional computed tomography study. *Eur J Orthod*, cjq190
- TOGASHI K, KITAURA H, YONETSU K, YOSHIDA N, NAKAMURA T. (2002) Three-dimensional cephalometry using helical computer tomography: measurement error caused by head inclination. *Angle Orthod*, 72, 513-520.
- TURHAN NA. (2009) Normal Okluzyona Sahip Ankara Bölgesi Çocuklarında Yaşa ve Cinsiyete göre sefalometrik normlar. Doktora

- tezi. Ankara Üniversitesi Sağlık Bilimleri Enstitüsü.
- TWEED, CH. (1946) The Francfort mandibular plane angle in orthodontics diagnosis, classification, treatment planning and prognosis. *Am. J. Orthod*, 32 :175-230
- UYSAI T, BAYSAL A, YAGCI A. (2009) Evaluation of speed, repeatability, and reproducibility of digital radiography with manual versus computer-assisted cephalometric analyses. *Eur J Orthod*, 31, 523-528.
- UZEL İ, ENACAR A. (2000) Ortodontide sefalometri, Baskı 2, Ankara. 3-210
- ÜÇÖK Ö, KAYADÜĞÜN A. (2015) Ortodontide Konik Işınli Bilgisayarlı Tomografi Endikasyonları (Bölüm 14). Güncel Bilgiler Işığında Ortodonti, E ÖZDİLER. 1. Baskı, Gümüş Kitapevi, Ankara, 283.
- ÜLGEN M, İŞCAN HN, ALTUĞ Z. (1982) Sefalometride Çizim ve Ölçüm Hataları (I); Aynı Bireylerin Belirli Zaman Aralıklarıyla Birbirinden Bağımsız Olarak Tekrarladıkları Çizim ve Ölçümleri Arasındaki Bireysel Farklılıklar. *Ankara Üniversitesi Dış Hekimliği Fakültesi Dergisi*, 9 (1) : 37 - 49.
- ÜLGEN M, ALTUĞ Z, İŞCAN HN. (1982) Sefalometride Çizim ve Ölçüm Hataları (II) ; Aynı Uzak Röntgen Resimlerinin Üç Araştırmacı Tarafından Yapılan Sefalometrik Ölçümleri Arasındaki Araştırmacılar Arası Farklılıklar. *Ankara Üniversitesi Dış Hekimliği Fakültesi Dergisi*, 9 (1): 77 - 89.
- ÜLGEN.M. (2000) Ortodontik anomaliler, sefalometri, etiyoloji, büyüme ve gelişim, tanı. Sayı 2. *İstanbul: T.C. Yeditepe Üniversitesi Yayınları*, 10-400.
- ÜNER O, İŞCAN HN, ILGAR S, AKKAYA S. (1986) Farklı baş konumlarında çekilen lateral sefalometrik filmlerde yumuşak dokuların incelenmesi. *G.Ü. Dışhek. Fak. Der. Cilt III, Sayı 2, Sayfa 21 - 35.*
- Van Der STELT PE. (2000) Principles of digital imaging. *Dent Clin North Am*, 44, 237-248.
- Van VLIJMEN OJ, BERGE SJ, SWENNEN GR, BRONKHORST EM, KATSAROS C, KUIJPERS-JAGTMAN AM. (2009) Comprasion of cephalometric radiographs obtained from cone-beam computed tomography scans and conventinel radiographs. *J Oral Maxillofac Surg*, 67:92-7
- Van VLIJMEN OJ, KUIJPERS MA, BERGE SJ, SCHOLS JG, MAAL TJ, BREUNING H, & KUILPERS-JAGTMAN AM. (2012) Evidence supporting the use of cone-beam computed tomography in orthodontics. *The Journal of the American Dental Association*, 143(3), 241-252.

- VAHDETTİN L, AKSOY S, ÖZ U, ORHAN K. (2016) Three-Dimensional cephalometric norms of Turkish Cypriots using CBCT images reconstructed from a volumetric rendering program in vivo. *Turkish Journal of Medical Scines*, 46(3), 848-861
- WHAITES E. (2006) Essentials of Dental Radiography and Radiology, 4th edition, Churchill Livingstone, UK,143–150
- WHITE S, PAE E. (2009) Patient image selection criteria for cone beam computed tomography imaging. *Semin Orthod*, 15, 19- 28.
- WU J, HÄGG U, RABIE ABM. (2007) Chinese norms of McNamara's cephalometric analysis. *Angle Orthod*, 77, 12–20.
- YAĞCI A, BÜYÜK SK. (2013) Mcnamara Cephalometric Norms Of Turkish Young Adults With Normal Occlusion And Well-Balanced Faces. *Journal of Health Sciences*, 22(1) 1-6.
- YOU KH, LEE KJ, LEE SH, BAIK HS. (2010) Three-dimensional computed tomography analysis of mandibular morphology in patients with facial asymmetry and man dibular prognathism. *Am J Orthod Dentofacial Orthop*, 138, 540.